

INTER-AMERICAN TROPICAL TUNA COMMISSION  
WORKING GROUP ON ECOSYSTEMS AND BYCATCH 3<sup>RD</sup>  
MEETING

La Jolla, California (USA) 26-27  
May 2025

**DOCUMENT EB-03 RD-F**

**Predicting albatross bycatch-risk hotspots across the North Pacific Ocean**

T. Clay, S. Shaffer, J. Adams, D. Anderson, S. Bograd, J. Felis, Y. Bedolla Guzmán, E. Hazen, R. Henry, M. Hester, D. Hyrenbach, J. Hernández Montoya, F. Jordan, F. Méndez Sánchez, B. Nishizawa, R. Orben, K. Ozaki, E. Rojas-Mayoral<sup>7</sup>, F. Sato, R. Suryan, J.-B. Thiebot, L. Thorne, H. Welch, L. Young, G. Carroll

**Abstract**

Bycatch threatens many seabird populations and remains a barrier to fisheries sustainability. Albatrosses are particularly vulnerable to bycatch in longline fisheries across the North Pacific due to their wide-ranging movements and attraction to fishing vessels. Identifying when, where and in which fisheries bycatch risk is greatest is crucial to prioritize monitoring and recommend targeted management interventions. Here, we collated >1,200 albatross tracks from eleven populations of three species (short-tailed, Laysan, and black-footed) in the North Pacific to provide an ocean-basin-scale assessment of bycatch risk. We overlaid species distribution model predictions of bird densities for each population and breeding stage with Automatic Identification Systems data on pelagic longline fishing effort and created indices of bycatch risk. Albatross distribution models had strong predictive power (mean [range] area under receiver operating characteristic curve: 0.91 [0.75-0.98]). We identified bycatch risk hotspots from pelagic longline fisheries in the central and northwest subtropical Pacific that occurred mostly during breeding (winter-spring); though the months with elevated risk differed according to population and fishing vessel flag state. By far the majority (98%) of overlap with pelagic longline fishing effort occurred within the WCPFC Convention Area, with only 2% occurring within the IATTC Convention Area. Three flag states (Japan, USA and Chinese Taipei) were responsible for >90% of total risk. Our results indicate that improved monitoring and increased adoption of and compliance with best-practice mitigation measures in high-risk fleets would reduce future conflicts between fisheries and albatrosses. Given the transboundary nature of seabird and fishing vessel movements, we also encourage harmonization of bycatch mitigation requirements between the IATTC and WCPFC.

**Introduction**

Bycatch is a major threat to seabirds, including North Pacific albatrosses – black-footed *Phoebastria nigripes*, Laysan *P. immutabilis* and short-tailed *P. albatrus* albatrosses – all of which are listed by the Agreement for the Conservation of Albatrosses and Petrels (ACAP; Lewison and Crowder, 2003; Phillips et al., 2016; Dias et al., 2019), while short-tailed albatrosses are also protected by the Endangered Species Act (ESA) in the United States (US). In the North Pacific, albatross bycatch rates were historically high in several fisheries, including US longline fleets, until the introduction of suite of mitigation measures in the early 2000s, which led to massive reductions in mortality, from tens of thousands to only hundreds of birds per year (Gilman et al., 2008; Melvin et al., 2011, 2019). Since the late 2000s, tuna Regional Fisheries Management Organizations (trFMOs) have mandated the use of best-practice mitigation in

pelagic longline fisheries, including use of streamer lines, night setting and line weighting through Conservation Management Measures (Western and Central Pacific Fisheries Commission [WCPFC]; CMM 2018-03; WCPFC 2018) or Resolutions (Inter-American Tropical Tuna Commission [IATTC] Resolution C-11-02; IATTC 2011). However, in many fleets, less than 5% of fishing sets are monitored, while many are not routinely monitored in all years (Phillips, 2013). Hence, there is often poor knowledge of where and when bycatch occurs, which fleets and vessels are responsible and whether they are appropriately using bycatch mitigation.

Given the logistical and political challenges and costs associated with collecting fisheries-dependent data, there is increasing emphasis on filling data gaps through alternative, fisheries-independent risk assessments (Clay et al., 2019; Good et al., 2023, 2024). Past research has shown that the three albatross species have wide-ranging distributions across the North Pacific and encounter and interact with a range of fisheries, predominantly pelagic and demersal longline and trawl (Hyrenbach and Dotson, 2003; Suryan et al., 2007; Fischer et al., 2009; Żydelis et al., 2011; Guy et al., 2013; Orben et al., 2021). To be able to sufficiently identify gaps in monitoring and develop targeted management interventions at the scale of RFMO areas of competency, there is need for standardized assessments of risk at ocean basin scales.

Here, we collate albatross tracking, oceanographic and fishing effort data to dynamically predict year-round distributions of North Pacific albatross populations and identify bycatch risk hotspots and the fishing fleets responsible. Specifically, we aimed to investigate 1) when and where albatross bycatch is most likely to occur, 2) the extent to which bycatch risk varies between the WCPFC and IATTC, and 3) the fleets and flag states responsible for the highest risk of interactions.

## Materials and Methods

### Tracking data collection and processing

All available tracking data were collated for the three North Pacific albatrosses and were requested through direct email contact with researchers or through the BirdLife International Seabird Tracking Database ([www.seabirdtracking.org](http://www.seabirdtracking.org)). We collated and standardized datasets from all tracked populations, breeding stages (incubation, brood, post-brood and non-breeding) and device types (GPS, PTT or geolocator) for the three species.

We plotted all GPS and PTT tracks and manually removed obviously erroneous locations. Geolocator tracks were estimated based on the timings of sunrise and sunset from light data using established routines, incorporating sea surface temperature (SST) information from tags with remotely-sensed maps to correct latitudes (for further details see Young et al., 2009; Jordan et al., 2022). Erroneous locations occurring two to three weeks either side of the spring and autumn equinox were removed. We took a conservative approach to further remove outliers beyond species' ranges which may lead to exaggerated bycatch risk estimates (e.g. in areas with high fishing intensity but low albatross usage). For black-footed and Laysan albatrosses, we calculated the 99% kernel utilization distribution (UD) isopleth based on all geolocator locations using the R package *adehabitatHR* (Calenge, 2006) and removed locations occurring outside this polygon.

We used the package *track2KBA* (Beal et al., 2021b) to split GPS and PTT tracks into discrete foraging trips, defined as periods of a minimum duration spent outside a spatial buffer around the breeding colony. For each population and breeding stage, we inspected the data and selected appropriate buffers that would exclude locations in the vicinity of the colony and remove incomplete trips. We used the package *foieGras* (Jonsen et al., 2020) to regularize tracks and remove erroneous locations based on a speed filter (90 km/hr). We selected an interpolated sampling interval of four hours for trips during breeding and an interval of twelve hours for short-tailed albatross PTT tracks, the latter to match the

temporal resolution of the non-breeding geolocator datasets.

### Species distribution models

Given that birds were tracked over a 20-year period (1998-2018), we built species distribution models (SDMs) to allow us to establish species-environment relationships and predict distributions across comparable time periods. We incorporated oceanographic variables known to influence albatross distributions as predictors in models (Hyrenbach et al., 2002; Suryan et al., 2006; Kappes et al., 2010; Żydelis et al., 2011). Daily dynamic environmental variables were acquired from Copernicus Marine Environmental Monitoring Service (<https://marine.copernicus.eu/>) and combined with topographic variables and distance from colony (for breeding data). See Welch et al. (2023) for details on variables and their sources.

We fit species distribution models using boosted regression trees (BRTs) due to their high predictive capability, fast computation time, and ability to fit non-linear relationships and handle collinearity among covariates. We generated pseudo-absences using random background selection at a ratio of 1:5 presences:absences. For breeding datasets, we randomly created absences for each colony within a buffer of the maximum range recorded for that species and breeding stage, which had been cropped by a land polygon to remove points occurring on land. For non-breeding datasets, we randomly created absences within a minimum convex polygon, which had been expanded by 200 km to account for geolocator error (Phillips et al., 2004) removing points occurring on land. Presence and pseudo-absence data were matched to environmental datasets in space and time, and BRTs with a binomial distribution were run for each species, population, and breeding stage. BRTs were built with a bag fraction of 0.6, a tree complexity of three and a learning rate between 0.0001 and 0.0001, to ensure that at least 2,000 trees were fit to each model.

Initial plotting of SDM outputs revealed that distance to colony had a strong effect for breeding models but that predictions were sensitive to the non-linear effect of distance from colony. To generate predictions that were more biologically meaningful and less sensitive to overfitting, we used the methodology of Hindell et al. (2020) and ran BRTs without distance variables. Instead, we created an accessibility layer for each population and breeding stage, and weighted model predictions for each location according to its accessibility from the breeding colony. Accessibility was modelled using a binomial distribution, in terms of the ratio of presences to absences, with distance to colony as a smoothed, monotonic decreasing constraint.

We evaluated the predictive performance of SDMs using the Area Under the Receiver Operator Characteristic Curve (AUC) metric, whereby values between 0.8 and 0.9, and 0.9 and 1.0, represent good and very good performance, respectively. We used two validation methods to assess performance: firstly, we trained models on a random subset of 70% of the data and tested it on the remaining 30%, iteratively 10 times (leave-out-70/30); secondly, for datasets for which we had two or more years of data, we trained on all years but one, and tested on the remaining year (leave-out-year), to test sensitivity of models to differences among tracking years.

Predictions were made for each day across a five-year period (2016-2020) corresponding to the availability of fishing effort data (below), within the spatial constraints of the training data. Daily predictions were averaged for each dataset and month and cropped using the threshold function in the *dismo* package (Hijmans et al., 2023) to remove “unsuitable” habitat. We weighted each probability of occurrence grid such that the sum of all values scaled to 1, to broadly match the structure of a utilization distribution (UD) grid.

### Weighting distributions according to phenology

We incorporated data on the timing of major events across the albatross annual cycle to weight model predictions and create monthly maps of predicted probability of presence at a population level, using an approach similar to Clay et al. (2019) and Carneiro et al. (2020). Data on the average timing of

1) return to the colony after non-breeding migration, 2) egg laying, 3) chick hatching, 4) end of chick brood and 5) departure of breeding adults from the colony, were obtained from the literature or from direct contact with data providers. As data were not available for all populations, we assumed phenology was the same for all populations of each species. The average timings were used to create weightings for each month, representing the proportion of that month covered by the four major breeding stages: incubation, brood, post-brood and non-breeding. For example, the average hatching date for Laysan albatrosses is 20 January and so the predicted probability of presence grid based on the incubation SDM was given a weighting of 0.65 (20 out of 31 days), while the probability of presence grid based on the brood SDM was given a weighting of 0.35 (11 out of 31 days). This process was repeated for each species, population and month. If a breeding stage was missing for a given population, that time period was not considered in analyses. We considered distributions during the pre-laying exodus period to be the same as during incubation (Clay et al., 2019).

### Fishing effort data and overlap analyses

We obtained Automated Identification System (AIS) fishing effort data v. 2 from Global Fishing Watch (<https://globalfishingwatch.org/data-download/datasets/public-fishing-effort>) for the period 2016-2020 at a 0.01° resolution. GFW combines public vessel registries and machine learning to identify fishing vessels and detect when they are actively fishing (Kroodsma et al. 2018). Data were resampled to a 2° resolution and effort (the number of fishing hours) in each cell was summed at a monthly level for each gear type and flag state.

We calculated monthly overlap per grid cell — an index of potential bycatch risk — as the predicted probability of albatross presence multiplied by the number of fishing hours (e.g. Jiménez et al., 2016; Clay et al., 2019; Carneiro et al., 2020), separately for each flag state. We also summarized overlap within Exclusive Economic Zones (EEZs; 200 nm from the coast) versus in the High Seas and within the Convention Areas of the IATTC versus the WCPFC. Monthly distribution grids were overlaid with shapefiles of country land borders and EEZs (v. 11; from <https://www.marineregions.org/downloads.php>). Here national jurisdictions are areas of EEZs each country including overseas dependencies and disputed territories (Beal et al., 2021a).

## **Results and Discussion**

Overall, we collated 1,243 tracks from eleven populations of the three species, including 428 tracks from four populations of black-footed albatrosses (at Torishima, Kure, Midway and Tern), 708 tracks from six populations of Laysan albatrosses (at Kure, Midway, Tern, Kaua'i, O'ahu and Guadalupe), and 107 track from short-tailed albatrosses at Torishima (Table 1). The populations represent around 51%, 78% and >85% of global breeding populations of the three species, respectively.

Through compiling the largest-ever database of albatross tracks in the North Pacific, building on previous studies (e.g. Suryan et al., 2007; Żydelis et al., 2011; Orben et al., 2021; Jordan et al., 2022) we show that the foraging distributions and exposure to fisheries are both species- and population-specific (Figure 1). Short-tailed albatrosses that breed in Japan forage along the Japanese coast and along the Aleutian Island chain in Alaska, Laysan albatrosses forage throughout the North Pacific and into the California Current, and black-footed albatrosses have a more sub-tropical distribution, but some populations forage in the northern California Current and Gulf of Alaska. The performance of species distribution models was very high for all species, especially for those based on GPS or PTT datasets (>0.96 for all species based on leave-out-70/30) (Table 2). Performance was lower for models based on geolocator data for Laysan and black-footed albatrosses (0.80 and 0.82 respectively) but still good. While AUC scores were slightly lower for leave-out-year validation, they were still very high for GPS and PTT datasets (>0.92 for all species) and moderate-to-high for geolocator datasets (0.80 and 0.77 for Laysan and black-footed albatrosses), indicating that SDMs still performed well when predicted onto different years of tracking data. As such, we conclude our models were suitable for predicting at-sea distributions and assessing coarse-scale risk outside the temporal range of the tracking data.

Across the North Pacific, we identified five bycatch risk hotspots for pelagic longline fisheries (Figure 2), many of which were shared among the three species. Risk was highest for black-footed and Laysan albatrosses 1) to the northeast of the main Hawaiian Islands and 2) to the northwest of the northwestern Hawaiian Islands, and for black-footed and short-tailed albatrosses 3) from Torishima up to the east coast of Honshu (Japan), and for Laysan albatrosses also 4) in a High Seas area of northwest Pacific east of Japan and 5) south of the main Hawaiian Islands. The majority (mean = 88%) of risk occurred in the High Seas, with values above 85% for black-footed and Laysan Albatross populations from the northwestern Hawaiian Islands (Kure, Midway and Tern) (Figure 3a). For both Torishima populations, most (78% and 99% for short-tailed and black-footed albatrosses, respectively) overlap with pelagic longline fisheries occurred within the EEZ of Japan, while for Laysan albatrosses from the main Hawaiian Islands (Kaua'i and O'ahu), a substantial proportion of overlap occurred within the US EEZ (85% and 27%, respectively).

Fleets operated by Japan, Chinese Taipei and the US contributed to 91% of total risk (Figure 4a). Broadly across the year, risk from pelagic longline fisheries was highest during the breeding season (November-May). Though, the specific months with elevated risk varied according to flag state (Figure 5a); there was elevated risk from US fleets during November to May and from the fleets Japan and Chinese Taipei during May to November, except for short-tailed albatrosses, for which risk was highest from Japanese fleets during October to May. There was also a distinct longitudinal pattern in bycatch risk, such that risk scores for western populations (Torishima, Kure and Midway) were more likely to be dominated by the fleets of Japan and Chinese Taipei, whereas risk scores for eastern populations (Tern, Kaua'i and O'ahu) were dominated by the US.

By far the majority (98%) of risk across all populations occurred within the WCPFC Convention Area west of 150°W, with only 2% occurring within the IATTC Convention Area (Figure 4b). Within the IATTC, 88% of total risk was from the US longline fleet and was east and northeast of the main Hawaiian Islands (Figure 2, 4b). The only population for which there was notable (11%) fisheries-overlap within the IATTC Convention Area was black-footed albatrosses from Tern Island, which overlapped mostly with US vessels (92%) during late breeding (March-June), and to a lesser extent with vessels flagged to Chinese Taipei (6%) during early breeding (November-December) (Figure 4a, b).

Overall, we show that most bycatch risk to North Pacific albatrosses from pelagic longline fisheries occurs within the WCPFC rather than the IATTC. In most high-risk areas identified, particularly in the northwest Pacific, fewer than 5% of vessels are monitored and use of best-practice bycatch mitigation (such as night setting, line weighting, use of streamer lines or hook-shielding devices) varies (Huang and Yeh, 2011; Sato et al., 2022). Additionally, given that US fisheries only contributed to around a third (32%) of overall risk, our findings suggest that recent bycatch reductions in US-managed fisheries (specifically in Hawai'i; Gilman et al., 2008) which have relatively high observer coverage (50-100%, depending on the fleet; e.g. NMFS 2023), may be insufficient to manage total risk for albatross populations, particularly northwestern populations. Fisheries overlap within the IATTC was mostly with the US shallow- and deep-set longline fleets, which have higher observer coverage and more stringent bycatch regulations than other fisheries in the North Pacific. However, the recent reduction in coverage of human observers in the US deep-set fishery from ~20% to 7% and the rapid roll-out of electronic monitoring may impact the ability to effectively monitor bycatch rates and mitigation use. We emphasize that increased adoption of best-practice mitigation measures and penalties for non-compliance across all fishing fleets would reduce future conflicts between fisheries and albatrosses. We also encourage the IATTC to update their bycatch mitigation requirements, so that they are harmonized with the WCPFC, given the transboundary nature of both seabird movements and fishing fleet dynamics. Lastly, given the importance of Guadalupe Island (Mexico) as a climate change refugia for Laysan albatrosses, efforts should be made to obtain more reliable fishing effort and bycatch data for fisheries in Mexican waters.

## Acknowledgements

We thank all the field workers and data managers, and funders that supported the collection, curation and analysis of the tracking data. In particular, we thank the Holocene Foundation for funding the data acquisition and analysis and are grateful to Global Fishing Watch for making their AIS fishing effort data openly available for use. We thank Steph Borrelle and Yasuko Suzuki for their useful comments on an earlier version.

## References

- Beal, M., Dias, M. P., Phillips, R. A., Opper, S., Hazin, C., Pearmain, E. J., et al. (2021a). Global political responsibility for the conservation of albatrosses and large petrels. *Sci. Adv.* 7, eabd7225. doi: 10.1126/sciadv.abd7225
- Beal, M., Opper, S., Handley, J., Pearmain, E. J., Morera-Pujol, V., Carneiro, A. P. B., et al. (2021b). track2KBA: An R package for identifying important sites for biodiversity from tracking data. *Methods Ecol. Evol.* n/a. doi: 10.1111/2041-210X.13713
- Calenge, C. (2006). The package “adehabitat” for the R software: A tool for the analysis of space and habitat use by animals. *Ecol. Model.* 197, 516–519. doi: 10.1016/j.ecolmodel.2006.03.017
- Carneiro, A. P. B., Pearmain, E. J., Opper, S., Clay, T. A., Phillips, R. A., Bonnet-Lebrun, A.-S., et al. (2020). A framework for mapping the distribution of seabirds by integrating tracking, demography and phenology. *J. Appl. Ecol.* n/a. doi: 10.1111/1365-2664.13568
- Clay, T. A., Small, C., Tuck, G. N., Pardo, D., Carneiro, A. P. B., Wood, A. G., et al. (2019). A comprehensive large-scale assessment of fisheries bycatch risk to threatened seabird populations. *J. Appl. Ecol.* 56, 1882–1893. doi: 10.1111/1365-2664.13407
- Dias, M. P., Martin, R., Pearmain, E. J., Burfield, I. J., Small, C., Phillips, R. A., et al. (2019). Threats to seabirds: A global assessment. *Biol. Conserv.* doi: 10.1016/j.biocon.2019.06.033
- Fischer, K. N., Suryan, R. M., Roby, D. D., and Balogh, G. R. (2009). Post-breeding season distribution of black-footed and Laysan albatrosses satellite-tagged in Alaska: Inter-specific differences in spatial overlap with North Pacific fisheries. *Biol. Conserv.* 142, 751–760. doi: 10.1016/j.biocon.2008.12.007
- Gilman, E., Kobayashi, D., and Chaloupka, M. (2008). Reducing seabird bycatch in the Hawaii longline tuna fishery. *Endanger. Species Res.* 5, 309–323. doi: 10.3354/esr00133
- Good, S. D., Dewar, K., Burns, P., Sainsbury, K., Phillips, R. A., Wallace, B. P., et al. (2024). Adapting the Marine Stewardship Council risk-based framework to estimate impacts on seabirds, marine mammals, marine turtles and sea snakes. *Mar. Policy* 163, 106118. doi: 10.1016/j.marpol.2024.106118
- Good, S. D., Gummery, M., McLennan, S., Dewar, K., Votier, S. C., and Phillips, R. A. (2023). Evaluating the appropriateness of risk-based approaches to assess the sustainability of fishery impacts on seabirds. *Endanger. Species Res.* 51, 161–172. doi: 10.3354/esr01251
- Guy, T. J., Jennings, S. L., Suryan, R. M., Melvin, E. F., Bellman, M. A., Ballance, L. T., et al. (2013). Overlap of North Pacific albatrosses with the U.S. west coast groundfish and shrimp fisheries. *Fish. Res.* 147, 222–234. doi: 10.1016/j.fishres.2013.06.009
- Hijmans, R. J., Phillips, S., and Elith, J. L. and J. (2023). *dismo: Species Distribution Modeling*. Available at: <https://cran.r-project.org/web/packages/dismo/index.html> (Accessed June 19, 2024).
- Hindell, M. A., Reisinger, R. R., Ropert-Coudert, Y., Hückstädt, L. A., Trathan, P. N., Bornemann, H., et al. (2020). Tracking of marine predators to protect Southern Ocean ecosystems. *Nature* 580, 87–92. doi:

10.1038/s41586-020-2126-y

Hyrenbach, K. D., and Dotson, R. C. (2003). Assessing the susceptibility of female black-footed albatross (*Phoebastria nigripes*) to longline fisheries during their post-breeding dispersal: an integrated approach. *Biol. Conserv.* 112, 391–404. doi: 10.1016/S0006-3207(02)00337-3

Hyrenbach, K. D., Fernández, P., and Anderson, D. J. (2002). Oceanographic habitats of two sympatric North Pacific albatrosses during the breeding season. *Mar. Ecol. Prog. Ser.* 233, 283–301. doi: 10.3354/meps233283

IATTC. 2011. Resolution C-11-02: Resolution to mitigate the impact on seabirds of fishing

Jiménez, S., Domingo, A., Brazeiro, A., Defeo, O., Wood, A. G., Froy, H., et al. (2016). Sex-related variation in the vulnerability of wandering albatrosses to pelagic longline fleets. *Anim. Conserv.* 19, 281–295. doi: 10.1111/acv.12245

Jonsen, I. D., Patterson, T. A., Costa, D. P., Doherty, P. D., Godley, B. J., Grecian, W. J., et al. (2020). A continuous-time state-space model for rapid quality control of argos locations from animal-borne tags. *Mov. Ecol.* 8, 31. doi: 10.1186/s40462-020-00217-7

Jordan, F. D., Shaffer, S. A., Connors, M. G., Stepanuk, J. E. F., Gilmour, M. E., Clatterbuck, C. A., et al. (2022). Divergent post-breeding spatial habitat use of Laysan and black-footed albatross. *Front. Ecol. Evol.* 10. doi: 10.3389/fevo.2022.1028317

Kappes, M. A., Shaffer, S. A., Tremblay, Y., Foley, D. G., Palacios, D. M., Robinson, P. W., et al. (2010). Hawaiian albatrosses track interannual variability of marine habitats in the North Pacific. *Prog. Oceanogr.* 86, 246–260. doi: 10.1016/j.pocean.2010.04.012

Lewison, R. L., and Crowder, L. B. (2003). Estimating Fishery Bycatch and Effects on a Vulnerable Seabird Population. *Ecol. Appl.* 13, 743–753. doi: 10.1890/1051-0761(2003)013[0743:EFBAEO]2.0.CO;2

Melvin, E. F., Dietrich, K. S., Fitzgerald, S., and Cardoso, T. (2011). Reducing seabird strikes with trawl cables in the pollock catcher-processor fleet in the eastern Bering Sea. *Polar Biol.* 34, 215–226. doi: 10.1007/s00300-010-0873-1

Melvin, E. F., Dietrich, K. S., Suryan, R. M., and Fitzgerald, S. M. (2019). Lessons from seabird conservation in Alaskan longline fisheries. *Conserv. Biol.* 33, 842–852. doi: 10.1111/cobi.13288

N.M.F.S. (2023). Seabird Interactions and Mitigation Efforts in Hawaii Pelagic Longline Fisheries 2022 Annual Report. Sustainable Fisheries Division, Pacific Islands regional Office, National Marine Fisheries Service. Issued September 2023.

Orben, R. A., Adams, J., Hester, M., Shaffer, S. A., Suryan, R. M., Deguchi, T., et al. (2021). Across borders: External factors and prior behaviour influence North Pacific albatross associations with fishing vessels. *J. Appl. Ecol.* 58, 1272–1283. doi: 10.1111/1365-2664.13849

Paolo, F., Kroodsma, D., Raynor, J., Hochberg, T., Davis, P., Cleary, J., et al. (2024). Satellite mapping reveals extensive industrial activity at sea. *Nature* 625, 85–91. doi: 10.1038/s41586-023-06825-8

Phillips, R. A. (2013). Requisite improvements to the estimation of seabird by-catch in pelagic longline fisherie. *Anim. Conserv.* doi: 10.1111/acv.12042

Phillips, R. A., Gales, R., Baker, G. B., Double, M. C., Favero, M., Quintana, F., et al. (2016). The conservation status and priorities for albatrosses and large petrels. *Biol. Conserv.* 201, 169–183. doi: 10.1016/j.biocon.2016.06.017

Phillips, R. A., Silk, J. R., Croxall, J. P., Afanasyev, V., and Briggs, D. R. (2004). Accuracy of geolocation estimates for flying seabirds. *Mar. Ecol. Prog. Ser.* 266, 265–272.

- Reynolds, M. H., Courtot, K. N., Berkowitz, P., Storlazzi, C. D., Moore, J., and Flint, E. (2015). Will the Effects of Sea-Level Rise Create Ecological Traps for Pacific Island Seabirds? *PLOS ONE* 10, e0136773. doi: 10.1371/journal.pone.0136773
- Suryan, R. M., Dietrich, K. S., Melvin, E. F., Balogh, G. R., Sato, F., and Ozaki, K. (2007). Migratory routes of short-tailed albatrosses: Use of exclusive economic zones of North Pacific Rim countries and spatial overlap with commercial fisheries in Alaska. *Biol. Conserv.* 137, 450–460. doi: 10.1016/j.biocon.2007.03.015
- Suryan, R. M., Sato, F., Balogh, G. R., David Hyrenbach, K., Sievert, P. R., and Ozaki, K. (2006). Foraging destinations and marine habitat use of short-tailed albatrosses: A multi-scale approach using first-passage time analysis. *Deep Sea Res. Part II Top. Stud. Oceanogr.* 53, 370–386. doi: 10.1016/j.dsr2.2006.01.012
- WCPFC. 2018. Conservation and Management Measure 2018-03: Conservation and Management Measure to Mitigate the Impact of Fishing for Highly Migratory Fish Stocks on Seabirds. Western and Central Pacific Fisheries Commission.
- Welch, H., Savoca, M. S., Brodie, S., Jacox, M. G., Muhling, B. A., Clay, T. A., et al. (2023). Impacts of marine heatwaves on top predator distributions are variable but predictable. *Nat. Commun.* 14, 5188. doi: 10.1038/s41467-023-40849-y
- Young, L. C., Vanderlip, C., Duffy, D. C., Afanasyev, V., and Shaffer, S. A. (2009). Bringing Home the Trash: Do Colony-Based Differences in Foraging Distribution Lead to Increased Plastic Ingestion in Laysan Albatrosses? *PLOS ONE* 4, e7623. doi: 10.1371/journal.pone.0007623
- Žydelis, R., Lewison, R. L., Shaffer, S. A., Moore, J. E., Boustany, A. M., Roberts, J. J., et al. (2011). Dynamic habitat models: using telemetry data to project fisheries bycatch. *Proc Biol Sci* 278, 3191–200. doi: 10.1098/rspb.2011.0330


**Table 1.** Summary of tracking data according to species and population.

Species	Region	Population	% global pop.	Breed stages	N. tracks
Laysan albatross	NW Hawaiian Is.	Kure (Mokupāpapa)	4.6	All	63
		Midway (Pihemanu)	72.8	All	105
		Tern (Kānemiloha‘i)	0.4	All	190
	Main Hawaiian Is.	Kaua‘i (Waipake & Nai Aina Kai)	<0.1	All	192
		O‘ahu	<0.1	Post-brood	84
Mexico Offshore Is.	Guadalupe	0.2	All	266	
Black-footed albatross	Japan Offshore Is.	Torishima	2.9	Incubation, brood, post-brood	21
	NW Hawaiian Is.	Kure (Mokupāpapa)	4.9	Brood, post-brood	158
		Midway (Pihemanu)	36.5	Post-brood	79
		Tern (Kānemiloha‘i)	7.1	All	170
Short-tailed albatross	Japan Offshore Is.	Torishima	85%	All	107


“All” under the Breed stage column represents that there were sufficient data for incubating, chick-rearing and non-breeding birds.

**Table 2.** Summary of species distribution model performance based on the Area Under the Receiver Operator Curve (AUC) statistic for each species, device type and model validation method. The median AUC score provided for each method with the range in parentheses. Leave-out-70/30 indicates where the model was trained on a random subset of 70% of the data and tested on the remaining 30%; Leave-out-year represents where the model was iteratively trained on all but one year of data and tested on the remaining year.


Validation method	Species	N datasets	AUC score (performance)	
			GPS/PTT	Geolocator
Leave-out-70/30	Laysan albatross	24	0.97 (0.94-0.99)	0.80 (0.68-0.86)
	Black-footed albatross	11	0.98 (0.95-0.99)	0.82 (0.80-0.84)
	Short-tailed albatross	4	0.96 (0.91-0.99)	–
Leave-out-year	Laysan albatross	10	0.95 (0.89-0.98)	0.80 (0.79-0.82)
	Black-footed albatross	6	0.95 (0.91-0.97)	0.77 (0.75-0.78)
	Short-tailed albatross	2	0.92 (0.88-0.97)	–




**Figure 1.** Compilation of more than 1,200 foraging tracks from 11 populations of three albatross species in the North Pacific Ocean. Blue tracks represent Laysan albatrosses, red tracks represent black-footed albatrosses and yellow tracks represent short-tailed albatrosses.




**Figure 2.** Bycatch-risk hotspots for short-tailed (top row), black-footed (middle row) and Laysan (bottom row) albatrosses in the North Pacific Ocean with pelagic longline fisheries. The color gradient indicates increasing risk (red represents the highest risk and light yellow represents the lowest risk) and hotspot areas are shown with black circles. The dark blue dashed line represents the boundary of the Convention Areas of the Inter-American Tropical Tuna Commission (IATTC) and the Western Central Pacific Fisheries Commission (WCPFC).



**Figure 3.** Bycatch risk from pelagic longline fisheries according to jurisdiction (Exclusive Economic Zone [EEZ] or High Seas) and population (bar). The average bycatch risk across the year is shown for each species and population, with westernmost populations on the left and easternmost populations on the right. Bars are filled according to jurisdiction, with the percentage of risk within the High Seas for each population specifically labeled at the bottom of each bar. STAL = short-tailed albatross; BFAL = black-footed albatross; LAAL = Laysan albatross; Tori = Torishima Island (Japan), Guad = Isla Guadalupe (Mexico).



**Figure 4.** Bycatch risk according to RFMO and flag state for each population. a) The average bycatch risk across the year is shown for each species and population, with westernmost populations on the left and easternmost populations on the right. Bars are filled according to fishing vessel flag state. The percentage of total risk within the IATTC Convention Area is shown for black-footed albatrosses from Midway Atoll, along with the flag states contributing to that risk. B) Important (>1%) flag states contributing to bycatch risk in each RFMO. STAL = short-tailed albatross; BFAL = black-footed albatross; LAAL = Laysan albatross; Tori = Torishima. Flag states: CHN = China; JPN = Japan; KOR = South Korea; Other = all other flag states combined; TWN = Chinese Taipei; USA = United States of America; VUT = Vanuatu.



**Figure 5.** Bycatch risk according to species, population, month and flag state a) across the entire North Pacific and b) just for black-footed albatrosses from Midway Atoll and Tern Island within the IATTC Convention Area. For each species (column) and population (row), the total bycatch risk in each month is shown, with the contribution of each flag state shown by a separate colour. Populations are ordered from the westernmost (top row) to the easternmost (bottom row) for each species.