

**INTER-AMERICAN TROPICAL TUNA COMMISSION**  
**AD HOC PERMANENT WORKING GROUP ON FADs**  
**9<sup>TH</sup> MEETING**  
**La Jolla, California (USA)**  
**28 - 29 May 2025**

**DOCUMENT FAD-09 INF-A**

**ANALYSES OF THE REGIONAL DATABASE OF STRANDED DRIFTING FISH  
AGGREGATING DEVICES (DFADS) IN THE PACIFIC OCEAN: A 2024  
UPDATE**

Mourot J.<sup>1</sup>, Thellier T.<sup>2</sup>, Lopez J.<sup>3</sup>, Fuller L.<sup>3</sup>, Román, M.<sup>3</sup>, David D.<sup>1</sup>, Ochavillo D.<sup>4</sup>, Smith W.<sup>5</sup>, Nicholas T.R.<sup>6</sup>, Tibatt B.<sup>7</sup>, Stevens K.<sup>8</sup>, Vaipuna L.<sup>9</sup>, Bigler B.<sup>10</sup>, Prioul F.<sup>11</sup>, Lercari M.<sup>11</sup>, Pollock K.<sup>12</sup>, Mesebeluu K.<sup>13</sup>, Ah Fook S.<sup>14</sup>, Iakopo M.<sup>14</sup>, Mesepitu J.<sup>15</sup>, Halumwane C.<sup>15</sup>, Batty M.<sup>16</sup>, Doutreloux N.<sup>17</sup>, Mugneret B.<sup>17</sup>, Lynch J.M.<sup>8</sup>, Tait H.<sup>5</sup>, Hamer P.<sup>1</sup> Escalle L.<sup>1</sup>

<sup>1</sup>The Pacific Community (SPC), Nouvelle-Calédonie

<sup>2</sup>Direction des ressources marines (DRM), Polynésie Française

<sup>3</sup>Inter-American Tropical Tuna Commission (IATTC)

<sup>4</sup>Department of Marine and Wildlife Resources (DMWR), American Samoa

<sup>5</sup>Tangaroa Blue Foundation (TBF), Australia

<sup>6</sup>Offshore Fisheries Division, Ministry of Marine Resources (MMR), Cook Islands

<sup>7</sup>Guam Department of Agriculture, Division of Aquatic and Wildlife Resources (DAWR), Guam

<sup>8</sup>Hawai'i Pacific University (HPU), Center for Marine Debris Research (CMDR), Hawai'i

<sup>9</sup>Ministry of Fisheries (MoF), Kingdom of Tonga

<sup>10</sup>Marshall Islands Marine Resources Authority (MIRMA), Marshall Islands

<sup>11</sup>Service du parc naturel de la mer de Corail et de la pêche (SPNMCP), Nouvelle-Calédonie

<sup>12</sup>Palmyra Atoll Program, The Nature Conservancy (TNC)

<sup>13</sup>Ministry of Agriculture, Fisheries, and the Environment (MAFE), Bureau of Fisheries, Republic of Palau

<sup>14</sup>Ministry of Agriculture and Fisheries (MAF), Samoa

<sup>15</sup>Ministry of Fisheries and Marine Resources (MFMR), Solomon Islands

<sup>16</sup>Tuvalu Fisheries Department, Tuvalu

<sup>17</sup>Direction des Services de l'Agriculture, de la Forêt et de la Pêche (DSA), Service de la Pêche et de Gestion des Ressources Marines, Wallis-and-Futuna

## SUMMARY

Drifting Fish Aggregating Devices (dFADs) are reaching coastal areas where they can become stranded, adding to pollution and/or causing environmental damage. To quantify these events and their impacts, several Pacific Island Countries and Territories (PICTs), in collaboration with the Pacific Community (SPC), and often with support from international Non-Governmental Organisations (NGOs), have implemented voluntary programmes to collect in-situ data. These data collection programs on stranded and lost dFADs are now fully implemented in sixteen PICTs: American Samoa, Australia, Cook Islands, Federated States of Micronesia, French Polynesia, Guam, Hawai'i, Republic of the Marshall Islands, New Caledonia, Palau, Palmyra, Solomon Islands, Tonga, Tuvalu, Samoa and Wallis and Futuna, with data collection spanning 2006–2025. This document provides an update of [FAD-07 INF-A](#) (Escalle et al., 2022) presented at the *Ad-hoc* Permanent Working Group on FADs (FADWG) in 2022. A total of 3,591 stranding events could be identified to date; 43.8% of these consisted of a buoy alone, 30.7% of a FAD alone and 21.6% of a FAD with a buoy attached (3.9% were unknowns). FADs and buoys were most commonly found on a beach (37.2%), while others had been previously collected by local communities (32.3%), and some were found drifting in the ocean (6.7%), or caught on coral reefs (4.3%). In some cases, it was possible to record environmental damages caused by dFAD strandings; this was most common for dFADs with submerged appendages and corresponded to coral damage (3.1% of all dFADs but 7.3% of all appendages found) or, very rarely, entanglements with animals (0.7% of all recorded dFADs and 0.8% of appendages found). Despite Pacific-wide studies have been recommended by the IATTC FADWG and supported by both the Western and Central Pacific Fisheries Commission (WCPFC) and the Inter-American Tropical Tuna Commission (IATTC), with Memorandums of Understanding (MoUs) on data sharing signed by the two organisations, it is important to note that no confidential information has been shared between organization to date. The origins of the stranded dFADs and buoys were investigated using markings on the buoys and satellite buoy serial numbers. Markings were compared with the public access vessel registry of the IATTC and WCPFC; while buoy serial numbers were matched with records in the IATTC and WCPFC observer data and the Parties to the Nauru Agreement (PNA) FAD tracking data. Stranded dFADs were coming from slightly more vessels fishing in the IATTC Convention Area (CA) (47.0%) than fishing in the WCPFC CA (34.1%), and 19.0% from vessels fishing in both CAs. Large variability in terms of country of origin for stranding events was observed. For example, most stranding events in French Polynesia originated from vessels fishing in the IATTC CA.

Generally, it should be noted that the data collection programme presented here provides an incomplete picture, highly dependent on data collection effect, of the level of dFAD strandings on Pacific Islands, and we suggest that additional countries and territories consider implementing similar data collection programs and participating in this regional initiative. Greater coverage of the dFAD stranding data is important to better understand the extent and potential implications of this issue and to help inform dFAD management options in the Pacific Ocean. Consequently, similar efforts and discussions are underway in the eastern Pacific Ocean to initiate voluntary regional data collection programmes on stranding dFADs to harmonize with the ongoing efforts in the Western and Central Pacific Ocean.

### 1. Introduction

Concerns regarding the number of drifting Fish Aggregating Devices (dFADs) reaching coastal waters and becoming stranded on sensitive habitats have been raised by several Pacific Island Countries and Territories (PICTs), regional entities and international Non-Governmental Organisations (NGOs). These concerns include the potential for dFADs to damage habitats such as coral reefs, entanglement of wildlife such as turtles and sharks, and contribute to coastline debris when stranded (Balderson and Martin, 2015; Escalle et al., 2019).

Such concerns have intensified in recent years due to a general perception of an increasing trend in stranding events, including in PICTs with no purse-seine activities in their EEZs, and by a lack of

retrieval plans and solutions to process/recycle the dFAD materials on remote islands. However, the number of studies investigating stranding events in the Pacific remains limited. This is largely due to the absence of data available to adequately quantify the number of dFADs arriving in coastal areas, stranding events, and impacts on ecosystems. Studies based on trajectories from satellite buoys deployed on dFADs operating in the Western and Central Pacific Ocean (WCPO) estimated that 11.3% of dFADs end up stranded (Escalle et al., 2023). However, the number of stranding events and level of ecosystem impacts are very likely under-estimated, given that the current dataset corresponds mostly to data from Parties of the Nauru Agreement (PNA) member EEZs, but also because satellite buoys are commonly deactivated by fishers when they drift outside their main fishing areas. To date, estimates of stranding events are also lacking in the eastern Pacific Ocean (EPO), although efforts to initiate regional data collection programs are underway.

This document — an update of [FAD-07 INF-A](#) — presents initiatives that have expended in the Western and Central Pacific Ocean and led PICTs in collaboration with the Pacific Community (SPC), local organisations, and/or NGOs, to collect data on lost dFADs reaching coastal waters and/or becoming stranded, as well as the impacts of these events on ecosystems. Data collection is carried out in each PICT and stored individually. These individual datasets are then compiled by SPC into a regional database with data from all PICTs, allowing for regional scientific studies to be performed, as well as the ground-truthing of existing estimates. An expansion of such data collection effort to the EPO, which is currently underway, would facilitate Pacific-wide analyses and an improved understanding of the impact of dFADs on the ecosystem. Consequently, supporting the initiation of regional data collection efforts in the EPO is also desirable.

## **2. Regional stranded FAD data collection programme**

Data collection programmes have been in place as early as 2004 (Australia), although several PICTs have either recently started or are developing programmes as a collaboration between SPC, national fisheries departments, local organisations and/or NGOs. These programmes collect data on recorded arrival events of dFADs in coastal areas and also address the need to collect in-situ data. Data collection programmes are in place in sixteen PICTs: American Samoa, Australia, Cook Islands, the Federated States of Micronesia, French Polynesia, Guam, Hawai'i, Kingdom of Tonga, New Caledonia, Palmyra, Republic of the Marshall Islands, Republic of Palau, Solomon Islands, Tuvalu, Samoa and Wallis and Futuna.

The main objectives of the programmes are to:

- quantify the number of dFAD stranding events or dFADs drifting nearshore;
- assess the resulting pollution and ecosystem impacts, including on species of special interest (SSIs) and key habitats;
- evaluate materials and designs of dFADs found stranded, in relation to past and current use of dFADs in the Pacific Ocean;
- evaluate how communities and PICTs may repurpose or recycle dFAD materials and satellite buoys locally, when possible;
- consider ways to mitigate the impacts of dFADs and provide scientific-based advice to guide the management of dFADs in the Pacific Ocean.

**TABLE 1.** Summary of data collected through stranded dFAD data collection programs in the Pacific Ocean.

PICT	Start of the program	Events recorded
French Polynesia	2019	1,539
Australia	2004	393
Cook Islands	2020	310
Wallis and Futuna	2020	268
Kingdom of Tonga	2023	201
Federated States of Micronesia	2021	187
Hawai'i (US)	2014	127
New Caledonia	2022	103
Republic of the Marshall Islands	2021	102
Solomon Islands	2024	93
Palmyra (US)	2009	86
Tuvalu	2022	61
Samoa	2024	28
American Samoa	2024	21
Pitcairn	Opportunistically	21
Vanuatu	Opportunistically	20
Guam	2024	8
Republic of Palau	2024	8
Wake Island (US)	Opportunistically	8
Papua New Guinea	Opportunistically/under discussion	4
Fiji	Opportunistically	2
New Zealand	Opportunistically	2
Alaska (US)	Opportunistically	1
Northern Mariana Islands (US)	Opportunistically	1
<b>Total</b>		<b>3,591</b>

Since 2020, data collection programmes have been developed by SPC in partnership with local fisheries departments and have started in the American Samoa; Cook Islands; the Federated States of Micronesia; Guam; New Caledonia; Republic of the Marshall Islands; Solomon Islands; Kingdom of Tonga; Tuvalu; Republic of Palau; Samoa and Wallis and Futuna (TABLE 1). Opportunistic data collection has also been reported to SPC since 2018, including through SPC's existing data collection networks, and includes additional records from Alaska, Fiji; New Zealand, Northern Mariana Islands, Papua New Guinea; Pitcairn Islands; Vanuatu; and Wake Island. These programs involve local communities reporting their findings to fisheries officers, who enter data on forms and in their country/territory database. Communication, dissemination, and awareness activities are essential because each programme depends on engagement by local communities. The types of awareness activities vary but can include posters, radio and TV broadcasts, and public talks (Appendix 1). For the first few months of the program in each PICT, reports included dFADs and buoys previously collected by the public. This information is important for creating a baseline inventory and for capturing and identifying new events. Data were also collected through dedicated visits to outer islands by SPC staff, national fisheries departments, and local staff (e.g., fisheries observers or fisheries officers). Island coastlines were then surveyed on a specific day, and data were collected for every dFAD found.

In parallel, other initiatives or opportunistic reports have emerged. This includes data collection at Palmyra Atoll since 2009 (through The Nature Conservancy; TNC); Hawai'i since 2014 (through the Center for Marine Debris Research); French Polynesia since 2019 (Marine Resources Authority); Australia since 2004 (Tangaroa Blue Foundation); and, very recently, Galapagos (Galapagos Conservation Trust as well as a private initiative conducted by the Tuna Conservation Group (TUNACONS) (TABLE 1). A description of data collection of stranded dFADs in these independent initiatives are detailed in

Appendix 2. Data were added to the regional database and analysed in this document, excluding data from Galapagos as none have been received by SPC.

Data fields collected by the PICTs include date, location, environment, materials, size and fate of the dFAD (e.g., removed, left where it was found, fished), the buoy identification alphanumeric code and any other painted marks on the buoy (often vessel names), as well as any observed environmental impacts (e.g., coral reef damage or entanglement of SSI). Data are collected in dedicated forms (Appendix 3), with details of each field in the form can be found in Appendix 4. Data are then transferred to SPC, where all of the data are compiled into a regional database.

### 3. Preliminary analyses

#### a) Summary of stranded events


A total of 3,591 stranding events were identified during 2006–2025 from all PICTs considered. Most of the stranding events consisted of buoys (43.8%), followed by FADs alone (30.7%), and by a FAD with a buoy attached (21.6%) (TABLE 2). FADs were either dFADs, including biodegradable dFADs (bio-FAD); or anchored FADs (aFADs), such as large metal drums used by some purse seine fleets in the WCPO (FIGURE 1 and FIGURE 2). The remaining events corresponded to a few radio buoys, and oceanographic buoys (FIGURE 1 and FIGURE 2), as well as a proportion of data where the type of floating object was not recorded (6.1%).

**TABLE 2.** Number of objects found stranded per type, with percentages of total in parentheses. The findings of FADs included Bio-FAD, dFADs, aFADs and dFAD parts (e.g., float, bamboo, or net found alone); buoys include satellite buoys, radio buoys, and oceanographic buoys.


		FAD (1,904)		
		Absence	Unknown	Presence
Buoys (2,448)	Absence	0	2 (0.1%)	1,103 (30.7%)
	Unknown	4 (0.1%)	10 (0.3%)	24 (0.7%)
	Presence	1,573 (43.8%)	98 (2.7%)	777 (21.6%)

The number of stranding events recorded in the regional database has been increasing with the development of data collection programmes and the growing number of PICTs participating. The first stranding events recorded were in 2006 in Australia with the launch of the Australian Marine Debris Initiative (AMDI) Database in 2004, followed by some records in Palmyra in 2009 (FIGURE 1). Data collection and awareness activities have been expanding since 2016, resulting in a gradual increase in the number of stranding events reported. As a result, more than 1,000 stranding events were reported in 2022, 447 in 2023 and 862 in 2024. In many countries, the first stage of the data collection program included an inventory of all buoys and FADs previously collected by local communities and often

accumulating in private properties or ports. Hence, the date is sometimes uncertain (11.6% of all stranding events) or unknown (0.6% of all stranding events). The dominant types of floating objects found in stranding events included satellite buoys, dFADs, and dFADs with a satellite buoy attached (FIGURE 1). The type of floating objects found stranded were slightly different between PICTs (FIGURE 2). For instance, in the Federated States of Micronesia and the Marshall Islands, respectively, 21.4% and 21.6% of the stranding events were industrial aFADs (FIGURE 2).



**FIGURE 1.** The number of stranded events found by year and type of FAD or buoy. The numbers at the top of the figure correspond to the number of stranding events per year.



**FIGURE 2.** Percentages of stranding events found by country and type of materials. Numbers at the top of the figure correspond to the number of stranding events per country. AK = Alaska; AS = American Samoa; AU = Australia; CK = Cook Islands; FJ = Fiji; FM = Federated States of Micronesia; GU = Guam; HW = Hawai'i; MH = Marshall Islands; MP = Northern Mariana Islands; NC = New Caledonia; NZ = New Zealand; PF = French Polynesia; PG = Papua New Guinea; PN = Pitcairn; PW = Palau; PY = Palmyra; SB = Solomon Islands; TO = Tonga; TV = Tuvalu; VU = Vanuatu; WF = Wallis and Futuna; WK = Wake Atoll; WS = Samoa.


Most of the buoys found were one of the three following brands: Satlink (44.3%), Marine Instruments (28.7%), and Zunibal (14.3%), and some were Ryokusei and Kato buoys (TABLE 3). Note that the brand was unknown for 10.7% of the buoys. Small differences between PICTs were detected (e.g., a higher proportion of Kato buoys was found in the Federated States of Micronesia).

**TABLE 3.** Brand of satellite buoys found stranded.


	Buoy brand	
	Number	%
Satlink	977	44.3
Marine Instruments	634	28.7
Zunibal	316	14.3
Kato	32	1.4
Ryokusei	11	0.5
Unknown	237	10.7
Total	2,207	

**b) Spatial distribution of stranding events**

The spatial distribution of FAD stranding events in the Pacific Ocean shows a large distribution over the PICTs where the data collection program is implemented (c 3 and FIGURE 4). A higher number of stranding events per 1° cell was detected in French Polynesia, Wallis and Futuna, one location in Solomon Islands and Tonga. However, it should be noted that this could be due to greater data collection efforts in those particular locations rather than a true reflection of higher levels of stranding events. Additional years of data and/or accounting for the effort in data collection is needed to better understand the spatial differences detected. In particular, in Solomon Islands, the data collection has only started in 2024 in specific locations and data collection in other locations might lead to similar level of stranding events.



**c 3.** Aggregated map of FADs found in Pacific Island Countries and Territories between 2006–2025. The legend represents the numbers of stranding events per 1° cells.



**FIGURE 4.** Map of stranding events with known positions (3,527) by type of object found in Pacific Island Countries and Territories and Australia between 2006–2025.

As shown in FIGURE 2, the type of floating objects was different depending on the country considered. For instance, the Federated States of Micronesia and the Marshall Islands detected more industrial aFADs than in other countries, or Australia presented mostly stranding events related to a buoy alone (92.9%) compared to a FAD alone, a FAD with a buoy attached or any other categories (1.8%, 4.3% and 1% respectively). Additional maps provided as supplementary materials in Appendix 5 (Figures S5.1 to S5.4) show locations of stranded FADs and/or buoys (i.e., raft, buoy, both or an unknown object), recorded in some of the four PICTs with the highest number of stranding events. For several PICTs, stranding events were greater on one side of the coast, (e.g., the case for some islands in the Tuamotu (French Polynesia, Figure S5.1). In the case of the Tuamotu Islands, a larger number of stranding events were detected on the east coasts (e.g., Rangiroa, Fakarava, Raraka), but it should be noted that greater data collection efforts occurred in these regions. One interesting case is the atoll of Raroia, with stranding events detected in the lagoon and the coasts inside the lagoon, likely after entering the lagoon on the east side. Some islands, such as Palmyra Atoll have a high density of coral reefs around their coastlines, making them sensitive locations to stranding events.

### c) Habitats impacted

FAD stranding events can occur in sensitive environments such as coral reefs and therefore can pose a risk to marine life and habitats. Out of the 3,591 stranding events recorded (FADs or buoys), 37.2% were found on a beach, 6.7% were drifting in the ocean and 4.2% were found on coral reefs (TABLE 4). Some of the data collected corresponds to objects previously collected by local communities and recorded as found in gardens or private properties, accounting for 32.3% of the data.

Results differ slightly when the type of object is considered separately (i.e., FAD or buoy). Buoys were mostly found in private properties (category “previously collected” in TABLE 4) (35.2%), followed by beaches (34.7%), then unknown (13.3%) and finally all the other habitats (less than 7.8% each). Buoys were often dismantled to recover electronic materials. In contrast, FADs were mostly found on a beach (41.0%), in private properties (25.4%), on a shore (9.9%) and on coral reefs (6.2%). The aFADs were mostly found on a beach (47.2%), private property (18.0%), on a shore (10.1%) or on a coral reef (7.9%). Results for dFADs varied depending on the presence of submerged appendages (i.e., a tail). DFADs with submerged appendages were more often found in private properties (35.9%), on a beach


(27.1%), drifting in the ocean (19.2%) or stranded on coral reefs (10.2%) compared to dFADs without any appendages (respectively 12.3%, 56.5%, 2.8%, and 5%).

**TABLE 4.** Numbers and percentages (in parentheses) of stranding events by habitat type and FAD.

Environment	Total	FADs	Buoys	DFAD with tail**	DFAD without tail**	AFAD
Anchored	1 (0.03%)	1 (0.1%)	1 (0.05%)	NA	NA	1 (1.1%)
Beach	1,331 (37.2%)	720 (41.0%)	765 (34.7%)	167 (27.1%)	364 (56.5%)	42 (47.2%)
Coral reef	154 (4.3%)	109 (6.2%)	57 (2.6%)	63 (10.2%)	32 (5%)	7 (7.9%)
Drifting in the lagoon	45 (1.3%)	35 (2%)	24 (1.1%)	17 (2.8%)	7 (1.1%)	4 (4.5%)
Drifting in the ocean	240 (6.7%)	160 (9.1%)	173 (7.8%)	118 (19.2%)	18 (2.8%)	4 (4.5%)
Mangrove	11 (0.3%)	8 (0.5%)	4 (0.2%)	NA	2 (0.3%)	6 (6.7%)
Previously collected*	1,153 (32.3%)	446 (25.4%)	777 (35.2%)	221 (35.9%)	79 (12.3%)	16 (18.0%)
Shore	249 (7.0%)	174 (9.9%)	100 (4.5%)	21 (3.4%)	127 (19.7%)	9 (10.1%)
Unknown	391 (10.9%)	105 (6%)	306 (13.9%)	9 (1.5%)	15 (2.3%)	NA

\*Found in a private property (garden, wharf or landfill). \*\*The term “tail” refers to dFADs’ submerged appendages.

The type of environment where FADs and buoys were found differed depending on the PICT considered. FIGURE 5 shows that a large proportion of objects were previously collected by local communities, who transformed and recycled materials, especially for buoys in the Cook Islands (20.6%) and New Caledonia (14.0%), and for both buoys and FADs in the Solomon Islands (95.7%; 86.4%), Samoa (96%; 80.0%), Tonga (82.2%; 79.6%), the Federated States of Micronesia (53.7%; 22.6%), French Polynesia (52.7%; 25.6%), Marshall Islands (21.1%; 52.7%) and Wallis and Futuna (13.4%; 16.5%). New Caledonia (24.3%), Palmyra Atoll (22.2%), the Federated States of Micronesia (14.5%), Australia (12.5%), Wallis and Futuna (9.45%), and Hawai’i (8.3%) also presented higher rates of FADs stranded on coral reefs.



**FIGURE 5.** Percentages of stranded buoys (top), and FADs (bottom) by habitat type and country. Numbers at the top of the figure correspond to the number of stranding events for each country. AS = American Samoa; AU = Australia; CK = Cook Islands; FM = Federated States of Micronesia; GU = Guam; HW = Hawai’i; MH = Marshall Islands; NC = New Caledonia; PF = French Polynesia; PG = Papua New Guinea; PN = Pitcairn; PW = Palau; PY = Palmyra; SB = Solomon Islands; TO = Tonga; TV = Tuvalu; VU = Vanuatu; WF = Wallis and Futuna; WK = Wake Atoll; WS = Samoa.

**d) Type of FADs found stranded**

The type and structure of FADs (i.e., dFAD and aFAD) found stranded was investigated. Slightly more FADs were found without submerged appendages (41.2%) compared FADs with submerged appendages (35.3%). However, this information was not recorded for 23.4% of all FADs (TABLE 5).

The condition of the FADs when found was also investigated, although this information was mostly not recorded (35.7%). When it was recorded, most FADs were found mostly fallen apart (28.6%), mainly without submerged appendages (14.2% of all FADs); followed by intact (23.9%), and again mainly without submerged appendages (13.8% of all FADs); and finally classified as beginning to break up (11.7%) (TABLE 5).

**TABLE 5.** Number (N) and percentages of stranded FADs with submerged appendages (left). Percentages of FAD condition and number of FADs in parentheses (right).


	Submerged appendages		Condition			
	N	%	Intact	Beginning break	to Mostly apart	fallen Unknown
Present	621	35.3	8.0% (141)	3.0% (53)	7.8% (138)	16.4% (289)
Absent	725	41.2	13.8% (242)	7.6% (133)	14.2% (249)	5.7% (101)
Unknown	412	23.4	2.2% (38)	1.1% (20)	6.6% (116)	13.5% (238)
Total	1,758		23.9% (421)	11.7% (206)	28.6% (503)	35.7% (628)

Materials used in the construction of FADs (i.e., classified as synthetic, natural or a mix of synthetic and natural materials) were also investigated (TABLE 6). Materials were not recorded for 15.7% of FADs. The remaining FADs (including aFADs) were made with a mix of synthetic and natural materials for rafts and no attachments (i.e., no tails; 26.2%), followed by synthetic rafts with no attachments (10.5%), and mixed rafts with synthetic appendages (10.1%). For all FADs found stranded, none were found with natural submerged appendages. Completely natural FADs without submerged appendages represented only 1.3% of all stranding events.

**TABLE 6.** Percentages and numbers (in parentheses) of FADs with the raft and submerged appendages made of synthetic, mixed, or natural materials (including structure, flotation and covering materials).

	Raft			
	Synthetic	Mix	Natural	Unknown
Synthetic tail	5.7% (101)	10.1% (178)	0.1% (2)	5.1% (89)
Mixed tail	0.5% (9)	1% (17)	0.1% (1)	0.7% (13)
Natural tail	0	0	0	0
No tail	10.5% (185)	26.2% (460)	1.3% (23)	3.2% (57)
Unknown	10.6% (187)	7.9% (139)	1.2% (21)	15.7% (276)


Materials were also investigated separately for the raft's main structure, the raft covering, and the submerged appendages. Structure and flotation materials were examined for 1,134 FADs (FIGURE 6A). Most of the structure and flotation materials detected in stranded FADs were i) bamboo and plastic flotation (42%); ii) bamboo (31.9%); and iii) plastic flotation (18.7%) (FIGURE 6A).



**FIGURE 6.** Percentage of flotation and structural materials (A); and covering materials (B) for FADs found stranded and with materials recorded (64.5% and 68.8% respectively) by country. The numbers at the top of each figure correspond to the number of stranding events with materials recorded by country. “Bamboo”, includes bamboo and/or log. “Plastic flotation” materials include float, PVC tube, plastic drum, polystyrene and plastic foam. AS = American Samoa; AU = Australia; CK = Cook Islands; FM = Federated States of Micronesia; GU = Guam; HW = Hawai’i; MH = Marshall Islands; NC = New Caledonia; PF = French Polynesia; PG = Papua New Guinea; PN = Pitcairn; PW = Palau; PY = Palmyra; SB = Solomon Islands; TO = Tonga; TV = Tuvalu; VU = Vanuatu; WF = Wallis and Futuna; WK = Wake Atoll; WS = Samoa.


When considering differences by countries, it can be noted that in the Federated States of Micronesia and the Marshall Islands, many of the stranding events were aFADs, and therefore flotation materials were recorded as metal drums or fiberglass (respectively 63.9% and 43.1%) (FIGURE 6A). The remaining materials for FADs found in these countries were a mix of bamboo, metal drum, plastic flotation and fiberglass (36.1% to 56.9%). FADs’ raft covering were typically made of netting and/or rope (85.8%). A higher percentage of FADs with no covering were also detected in the Federated States of Micronesia and the Marshall Islands, mainly corresponding to aFADs (FIGURE 6B).

The presence or absence of FAD netting in the raft was not often recorded. When this information was recorded, 22.0% of FADs did not have any netting (mostly aFADs) and 16.7% of FADs had some netting as covering, but details about mesh size were not recorded (FIGURE 7). When mesh size was recorded, 43.7% of FADs had small mesh netting (<7cm), 22.7% had large mesh netting (≥7cm) and 11.7% had both small and large mesh netting (FIGURE 7).



**FIGURE 7.** The percentage of visually estimated mesh size used to cover the rafts of FADs (small: <7cm, large: ≥7cm; or a combination of small and large netting), when recorded (42.9% of unknown removed), found stranded by country. The numbers at the top of the figure correspond to the number of stranded events with materials recorded by country. AS = American Samoa; AU = Australia; CK = Cook Island; FM = Federated States of Micronesia; GU = Guam; HW = Hawai’i; MH = Marshall Islands; NC = New Caledonia; PF = French Polynesia; PN = Pitcairn; PW = Palau; PY = Palmyra; SB = Solomon Islands; TO = Tonga; TV = Tuvalu; WF = Wallis and Futuna; WK = Wake Atoll; WS = Samoa.

The most common materials used to construct the submerged appendages attached to FADs were netting and/or rope (76.2%). The remaining (23.8%) were constructed with a combination of bamboo, plastic materials, net, and weights (FIGURE 8). In New Caledonia, all the FADs found stranded with submerged appendages were composed of netting, which creates a high risk for coral entanglement. Despite the high numbers of submerged appendage materials recorded as unknown (34.4%), when netting was recorded, the mesh size, as well as the design, were also examined. Small mesh netting (<7 cm) was found in 45.5% of records, compared to 32.7% with large mesh netting (TABLE 7). Even if a large proportion presented no information on the design used (36.6%), most of the FADs found had an open panel (33.7%) followed closely by submerged appendages rolled up in a bundle (25.4%) (TABLE 7).




**FIGURE 8.** Materials used for the construction of submerged appendages of FADs found stranded, recorded by country. The numbers at the top of the figure correspond to the number of stranded events with materials recorded by country. Plastic materials include plastic sheeting, plastic drums, fishing line, PVC tubes and float. For reading clarity reasons, the category “Other” is a mix of categories which each represent a low number of stranding events (<4) (combination of bamboo, net, plastic materials, weight, metal and/or fabric). ”w/wth”= with or without. AS = American Samoa; AU = Australia; CK = Cook Islands; FM = Federated States of Micronesia; HW = Hawai’i; MH = Marshall

Islands; NC = New Caledonia; PF = French Polynesia; PG = Papua New Guinea; PN = Pitcairn; PY = Palmyra; SB = Solomon Islands; TO = Tonga; TV = Tuvalu; WF = Wallis and Futuna; WK = Wake Atoll; WS = Samoa.

**TABLE 7.** Design (left) and mesh size (right) of netting used as submerged appendages of stranded FADs.

Design	Numbers	Percentages	Mesh net size	Numbers	Percentages
Unknown design	111	36.6%	Small (< 7 cm)	138	45.5%
Open panel	102	33.7%	Large (≥7cm)	99	32.7%
Rolled up in a bundle	77	25.4%	Unknown size	36	11.9%
Mixed design	13	4.3%	Small and large	30	9.9%

In 44.6% of the stranding events, the shape of the FAD rafts was recorded, and different shapes of rafts were detected (FIGURE 9 and TABLE 8). Rectangular and square rafts were the most common (22.6% and 11.8%, respectively), followed by cylindrical rafts (4.6%).



**FIGURE 9.** Shape of FADs found stranded by country. Numbers at the top of the figure correspond to the number of stranded events with FAD shape recorded per country. AS = American Samoa; AU = Australia; CK = Cook Islands; FM = Federated States of Micronesia; GU = Guam; HW = Hawai'i; MH = Marshall Islands; NC = New Caledonia; PF = French Polynesia; PG = Papua New Guinea; PN = Pitcairn; PW = Palau; PY = Palmyra; SB = Solomon Islands; TO = Tonga; WF = Wallis and Futuna.

**TABLE 8.** Shapes of the FADs found stranded in all the Pacific Islands Countries and Territories.

Shape of the raft	Unknown	Rectangular	Square	Buoy sausage	Cylindrical	PVC Sausage	Boat shape	Octagonal
Percentage	55.4%	22.6%	11.8%	4.9%	4.6%	0.4%	0.2%	0.1%
(Number)	(974)	(398)	(207)	(86)	(81)	(7)	(3)	(2)

### e) Environmental impacts

The fate of buoys and FADs found stranded was investigated (TABLE 9). Most of the buoys (75.5%) were removed from the environment, while a lower number of FADs (28.0%) were removed. It should be noted that in a large portion of the stranding events, the fate was not recorded (18.6% of buoys and 51.8% of FADs).

**TABLE 9.** Fate of buoys and FADs found stranded.

	Buoy		FAD	
	Number	%	Number	%
Removed	1,849	75.5	534	28.0
Left	143	5.8	348	18.3
Removed partly	NA	NA	16	0.8
Sunk	NA	NA	8	0.4
Fished and left	NA	NA	6	0.3
Fished and removed	NA	NA	4	0.2
Relocated and left	NA	NA	2	0.1
Unknown	456	18.6	986	51.8

The purpose of the removal of buoys and FADs from the environment was recorded when possible (TABLE 10). This information was available for 59.8% of the buoys and 80.3% of the FADs. Buoys were mostly removed to be placed in storage (19.6%), in a landfill (10.7%), or left with the finder (8.1%). Communities also reused buoys (9.9%): using them as home furniture, like flowerpots (32.8%), as a light source (14.8%), using the electronic components such as solar panels or batteries (9.3%) or for fishing activities (2.1%). Most of the FADs removed from the environment were reused (48.3%) either transformed into house furniture (86.8%), or for fishing activities or boat furniture (4.7%). Some of the remaining FADs were used for research (9.6%), placed in a landfill (8.5%) or stored at the finder’s home (8.8%) or elsewhere (3.8%). It should be noted that fate and purpose of removed buoys and FADs was highly variable between PICTs and not necessarily classified the same way everywhere.

**TABLE 10.** Investigation of the purpose and fate of buoys and FADs removed from the environment.

Purpose	FADs		Buoys	
	Number	Percentage	Number	Percentage
Reused	257	48.3	183	9.9
Unknown	105	19.7	744	40.2
Research	51	9.6	46	2.5
Left with the finder	47	8.8	150	8.1
Landfill	45	8.5	197	10.7
Storage	20	3.8	363	19.6
Burned	4	0.8	NA	NA
Dismantled	3	0.6	11	0.6
Reused/Stored for ReCon	NA	NA	153	8.3
Relocated	NA	NA	2	0.1

Environmental damage could be recorded (36.8% of all FADs found stranded) and corresponded mostly to dFADs with submerged appendages. Damage was associated with coral (3.1% of all dFADs but 7.3% of all FADs with appendages found) or, very rarely, entanglements with animals (0.7% of all dFADs but 0.8% of all FADs with appendages found) (TABLE 11). Few FADs have been reported as entangled, however it was not precisely determined if it was on corals or on rocks, therefore it is mentioned as “unidentified” (1.7% of all FADs). It should be noted that the environmental damage was recorded at the time of locating a FAD. However, these may be underestimated, as ghost fishing, marine pollution or coral damage can potentially occur throughout the lifetime of FADs (at-sea or on

coastal habitats). Marine pollution can also occur through microplastics from FAD parts or heavy metal pollution from electronic components and batteries in satellite buoys, but this would be difficult to quantify under the current data collection methods.

**TABLE 11.** Environmental damage caused by stranded FADs recorded in the database.

	Total FADs	dFAD	dFAD with tail	dFAD without tail	aFAD
Entangled animals	12 (0.7%)	11 (0.7%)	5 (0.8%)	3 (0.5%)	1 (1.1%)
Entangled on corals	54 (3.1%)	51 (3.1%)	45 (7.3%)	1 (0.2%)	3 (3.4%)
Entangled on corals and animals	1 (0.1%)	1 (0.1%)	1 (0.2%)	NA	NA
Entangled (unidentified)	30 (1.7%)	30 (1.8%)	22 (3.6%)	7 (1.1%)	NA
No	592 (33.7%)	542 (32.5%)	139 (22.6%)	320 (49.7%)	50 (56.2%)
Unknown	1,069 (60.8%)	1034 (62%)	404 (65.6%)	313 (48.6%)	35 (39.3%)

Environmental damage, particularly related to coral, by the submerged appendages was also investigated. Few records of coral damage and unidentified entanglements were recorded (45 and 22 respectively). Most dFADs with submerged appendages found entangled on corals involved netting with small mesh size (TABLE 12 A). Most of dFADs with submerged appendages found entangled on corals also involved a design with open panels (29.2%) (TABLE 12 B). However, the net mesh size, or the design were often not recorded (for coral damage, respectively 24.6% and 26.2%; for unidentified damage, respectively 7.7% and 6.2%).

**TABLE 12.** Percentage and number (in parentheses) of FADs found with submerged appendages entangled on corals and unidentified, depending on the netting mesh size (A) or the design (B).

(A)	Small (<7cm)	Large (≥7cm)	Small and large	Unknown size
Coral damage	15.4% (10)	10.8% (7)	15.4% (10)	24.6% (16)
Unidentified damage	18.5% (12)	4.6% (3)	3.1% (2)	7.7% (5)

(B)	Open panel	Rolled up into a bundle	Mixed design	Unknown design
Coral damage	29.2% (19)	6.2% (4)	4.6% (3)	26.2% (17)
Unidentified damage	16.9% (11)	9.2% (6)	1.5% (1)	6.2% (4)

**f) Origin – Matching with observer and FAD tracking data**


Two approaches were used to determine the origin of the FADs and buoys found stranded in the Pacific Ocean. First, the marks painted on the buoys were used to identify the vessel monitoring the buoy. Marks on the satellite buoys were compared to the WCPFC and IATTC online vessel registry to identify the possible vessels, which allowed identification of flag and Convention Area (CA) where the

owner vessel has been fishing. The second approach used to determine the origin of the stranded buoys consisted of identifying the unique buoy ID alphanumeric codes from the database and cross referencing it against three fishery databases: i) the PNA FAD tracking database; ii) the WCPFC observer database; and iii) the IATTC observer database. The last known position in the PNA FAD tracking data and/or the last activity recorded in the observer data from WCPFC and IATTC was identified for each buoy that has a unique ID number that matched a number in the corresponding database. Despite Pacific-wide studies have been recommended by the IATTC FADWG and supported by both the WCPFC and the IATTC, with Memorandums of Understanding (MoUs) on data sharing signed by the two organisations, it is important to note that no confidential information has been shared between organization to date


A painted mark was found on 56.6% of the 2,207 satellite buoys found. 19.0% of the buoys did not have any marks and the presence of any marking was unknown for 24.4% of the buoys. However, markings on the buoy did not always result in the identification of a vessel or a flag. Among all the satellite buoys with painted marks, 32.4% could lead to the identification of the flag of the owner vessel and the CA where it has been fishing. Also to note that a few satellite buoys presented only a letter as a marking (14.3%), making it therefore impossible to identify a vessel, thus the flag or the CA.

A large variability in the origin of vessels monitoring satellite buoys found stranded (and attached to a dFAD or not) was detected (FIGURE 10). 29.1% of buoys were from Ecuadorian vessels; 23.2% from United States of America (US) vessels; 11.9% from Korean vessels; 10.9% from Panamean vessels and the rest from 17 other flags (FIGURE 10). Most buoys found stranded were from vessels fishing in the IATTC CA (47.0%), followed by WCPFC CA (34.1%) and both CAs (19.0%) (FIGURE 11A). With the second method using the unique buoy identification number, among the 2,207 stranded satellites buoys, 20.8% had been found in one of three fishery databases with 41.6% from the IATTC observer database, 36.4% from the PNA FAD Tracking database and 22.0% from the WCPFC observer database. However, the last known position investigated highlighted that 52.9% of buoys were last recorded in the WCPFC CA, 44.2% in the IATTC CA, and only 2.8% of them in the IATTC/WCPFC overlap area (see FIGURE 12 for the area covered by each CA and the overlap area).


In terms of country of stranding events, 86.4% of the marks on buoys found in French Polynesia were from vessels of the IATTC CA (Ecuador, Panama, US and Colombia) (FIGURE 10) and 85.1% of the last recorded positions were from the IATTC CA (FIGURE 11B). The buoys found with marks in the Federated States of Micronesia were almost exclusively from vessels fishing in the WCPFC CA (86.4%), although from a wide range of fleets: Korea, Japan, US, Nauru, Ecuador, Chinese Taipei and Papua New Guinea as examples (FIGURE 10 and FIGURE 11A). In Cook Islands, a large range of vessel flags from the IATTC CA were detected from marked buoys, mostly from US, Ecuadorian and Panamean vessels (36.1, 22.2%, and 11.1%, respectively). Finally, stranding events in Australia, located in the western part of the Pacific Ocean, consisted mostly of buoys marked from US (34.6%), Korea (25.0%), Colombia and Ecuador (5.8% each).



**FIGURE 10.** Flag of owner vessel identified using marks painted on the satellite buoys by stranding location, using publicly available IATTC and WCPFC vessel registers. Numbers at the top of the figure correspond to the number of buoys stranded per country. AS = American Samoa; AU = Australia; CK = Cook Islands; CN = China; CO = Columbia; EC = Ecuador; FM = Federated States of Micronesia; GU = Guam; HN = Honduras; HW = Hawai’i; JP = Japan; KI = Kiribati; KR = Korea; MH = Marshall Islands; MX = Mexico; NC = New Caledonia; NR = Nauru; PA = Panama; PE = Peru; PF = French Polynesia; PG = Papua New Guinea; PN = Pitcairn; PY = Palmyra; SB = Solomon Islands; TO = Tonga; TV = Tuvalu; TW = Chinese Taipei; US = USA; VU = Vanuatu; VZ = Venezuela; WF = Wallis and Futuna; WK = Wake Atoll; WS = Samoa.




**FIGURE 11.** Convention area of owner vessel identified using marks painted on the satellite buoys, using publicly available IATTC and WCPFC vessel registers (A) and convention area of the last known position in the PNA FAD tracking data, the WCPFC and IATTC observer data (noting no confidential information was shared; e.g. vessel ID, flag ID, satellite buoy track) (B) by stranded location. IATTC/WCPFC = vessel fishing in both convention areas (A) or the overlap area between IATTC and WCPFC (B). Numbers at the top of the figure correspond to the number of buoys stranded per country. AS = American Samoa; AU = Australia; CK = Cook Islands; FM = Federated States of Micronesia; HW = Hawai’i; MH = Marshall Islands; MP = Northern Mariana Islands; NC = New Caledonia; PF = French Polynesia; PN = Pitcairn; PY = Palmyra; SB = Solomon Islands; TO = Tonga; TV = Tuvalu; VU = Vanuatu; WF = Wallis and Futuna; WK = Wake Atoll; WS = Samoa.




**FIGURE 12.** Map of the WCPFC and IATTC Convention Areas, including the overlap area. Areas of international waters I1, I2, and I3, as used in FIGURE 16, are indicated in green, pink and light green.

The time difference between the last known date and the date found stranded was investigated using the second approach. This time difference was then calculated and categorized into three classes: less than one year; between one and two years; and more than two years. Maps indicating the last recorded positions of buoys found stranded, and time between the last known date and the date found stranded have been compiled (FIGURE 13 and FIGURE 14).


As previously identified, certain PICTs received satellite buoys mostly from one CA only (FIGURE 15). For instance, French Polynesia, Hawai'i, and Palmyra Atoll have stranded buoys mostly from vessels fishing in the IATTC CA and few from the WCPFC CA (FIGURE 15). It was also found that buoys were mainly drifting and stranded for more than one year or two years before being found stranded. Australia received buoys coming from WCPFC CA, which could be linked to geographical proximity, but there are also buoys coming from the IATTC CA, one and two years after the last recorded activity (FIGURE 15).




**FIGURE 13.** Map with all recorded stranded positions (black dots) and last known positions (green olive symbols) from buoys stranded and found in the IATTC observer data only (noting no confidential information was shared; e.g. vessel ID, flag ID, satellite buoy track). The color of the lines indicates the time between last known position and the date found stranded.



**FIGURE 14.** Map with all recorded stranded positions (black dots) and last known positions (pink symbols) from buoys stranded and found in the WCPFC observer data only (noting no confidential information was shared; e.g. vessel ID, flag ID, satellite buoy track). The color of the lines indicates the time between last known position and the date found stranded.





**FIGURE 15.** Maps of Hawai’i, Palmyra Atoll, Wake Atoll, Guam, Northern Mariana Islands (A), Cook Islands (B) and French Polynesia (C) with buoy stranded positions (black dots) and the last known position of buoys from three fishery databases: the PNA FAD tracking data (light green symbols); the WCPFC observer data (pink symbols) and the IATTC observer data (green olive symbols; noting no confidential information was shared; e.g. vessel ID, flag ID, satellite buoy track). The color of the lines indicates the time between the last known position and the date found stranded.

The exclusive economic zone (EEZ) of the last known position of buoys was investigated to detect potential connectivity and movement patterns of origin by stranding areas (FIGURE 16). In the following paragraph, a focus is made on the countries located in the western part of the Pacific Ocean. In the Cook Islands, most stranded FADs were last detected in the high seas in the eastern Pacific Ocean (13; 40%), the high seas in the central Pacific Ocean (12; 27.1%), the Kiribati Line Islands (15.7%) and the Galapagos (4.3 %) (FIGURE 16). In the Federated States of Micronesia, most stranded FADs were last detected in Federated States of Micronesia (56.4%), Kiribati Gilbert Islands (14.5%), Papua New Guinea (9.1%) and the high seas in the central Pacific Ocean (12; 5.5%).

Stranding areas	WCPFC CA																Origin areas										IATTC CA			
	AU	PG	FM	SB	MH	WK	VU	NR	GL	LN	PX	I1	TV	FJ	WF	TK	AS	CK	PY	HW	I2	I3	GP	PU	CR	CO	EC			
AS	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0			
AU	3.3	7.9	0	21.1	0	0	1.3	0	5.3	8.2	2.6	8.2	11.8	2.6	0	2.6	0	2.6	1.3	0	16.8	6.3	0	0	0	0	0			
CK	0	1.4	0	0	0	0	0	0	15.7	1.4	2.9	2.9	0	0	0	0	0	1.4	0	0	27.1	4.3	4.3	0	0	0	2.9			
FM	0	9.1	56.4	1.8	1.8	0	0	0	14.5	3.6	1.8	3.6	0	0	0	0	0	0	0	0	5.5	1.8	0	0	0	0	0			
GU	0	0	0	0	33.3	0	0	0	33.3	0	0	33.3	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0			
HW	0	0	0	0	9.1	0	0	0	9.1	9.1	0	0	0	0	0	0	0	0	0	9.1	83.6	0	0	0	0	0	0			
MH	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0			
MP	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0			
NC	0	0	0	19.4	0	0	0	0	12.9	5.5	5.5	9.7	3.2	0	3.2	0	0	0	0	12.9	16.1	9.7	0	0	0	0	0			
PF	0	1.1	0	0	0	0	0	1.1	1.1	2.1	4.3	1.1	0	0	0	0	0	0	0	30.9	49.7	6.4	1.1	1.1	2.1	2.1				
PY	0	0	0	0	0	0	0	0	15.7	0	16.7	0	0	0	0	0	0	0	16.7	0	33.3	16.7	0	0	0	0	0			
SB	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0			
TO	0	0	0	0	2.7	0	0	2.7	13.5	0	2.7	8.1	0	0	0	2.7	0	0	0	29.7	32.4	5.4	0	0	0	0	0			
TV	0	0	0	0	0	0	0	0	16.7	0	0	5.0	0	0	0	0	0	0	0	33.3	0	0	0	0	0	0	0			
VU	0	0	0	42.9	0	0	0	0	0	0	0	28.5	28.5	0	0	0	0	0	0	0	0	0	0	0	0	0	0			
WF	0	0	0	0	0	0	0	0	24.4	6.7	6.7	33.3	0	4.4	2.2	0	6.7	0	0	11.1	4.4	0	0	0	0	0	0			
WK	0	0	0	0	0	20	0	0	0	0	20	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0			
WS	0	0	0	0	0	0	0	0	5.0	0	0	0	0	0	0	0	0	0	0	5.0	0	0	0	0	0	0	0			


**FIGURE 16.** Matrix showing EEZ of origin for FADs found stranded: stranding country (left) and EEZ or origin (top), derived from the stranded position and the last known position in the PNA FAD tracking data, the WCPFC observer data or the IATTC observer data (noting no confidential information was shared; e.g. vessel ID, flag ID, satellite buoy track). AS = American Samoa; AU = Australia; CK = Cook Islands; CO = Colombia; CR = Costa Rica; EC = Ecuador; FJ = Fiji; FM = Federated States of Micronesia; GL = Gilbert Islands; GP = Galapagos; GU = Guam; HW = Hawai’i; LN = Line Islands; MH = Marshall Islands; MP = Northern Mariana Islands; NC = New Caledonia; NR = Nauru; PF = French Polynesia; PG = Papua New Guinea; PU = Peru; PY = Palmyra ; PX = Phoenix islands; SB = Solomon Islands; TK = Tokelau; TO = Tonga; TV = Tuvalu; VU = Vanuatu; WF = Wallis and Futuna; WK = Wake Atoll; WS = Samoa; I1 = Internal waters between Gilbert, Phoenix and Line Islands (174°–202°); I2 = International waters East of the Line Islands and North of French Polynesia (202°–240°) and I3 = Eastern part of the Pacific Ocean (east of 240°), see FIGURE 12.

In French Polynesia, most stranded FADs were last detected in the IATTC CA, in the high seas of the central and eastern part of the Pacific Ocean (I3 and I2, 45.7% and 30.9%, respectively) (FIGURE 16). In Hawai'i and Palmyra, most stranded dFADs were last detected in the IATTC CA in the high seas of the central part of the Pacific Ocean (I2) (63.6% and 33.3%, respectively). In Wallis and Futuna, most stranded dFADs were last detected in the WCPFC CA, in Tuvalu (33.3%) and the Line Islands (24.4%), as well as the IATTC high-seas areas in the central part of the Pacific Ocean (I2, 11.1%). Some countries such as American Samoa, Northern Mariana Islands and Solomon Islands, present 100% of last detections from I2 or I3 zones but a low number of buoys have been found in the different fishery databases for those 3 countries.

The time difference between the date buoys and FADs were found stranded and their last known position was investigated (FIGURE 17). 20.4% of all buoys were found less than one year after the time of their last known position, 27.4% were within two years, 38.1% of them were found more than years after the time of their last known position. For 14.2% of them the range of time is unknown.

In most PICTs, the data collection programs started recently but may recorded FADs and buoys found years ago by communities. In addition, data on stranding events were collected between 2006 and 2025. Hence, the range of years between the date found stranded and the last known position was highly variable in some PICTs. For example, it reached more than 4,000 days (about 11 years) for some buoys found in French Polynesia. It can also be noted that the time differences varied depending on the database used, for instance, higher time differences were detected for matches with the WCPFC observer's database, which recorded the last activity in the observer data; and smaller time differences for matchings with the PNA FAD tracking data (with available data starting in 2016), which is closer to the real date of last transmission. In the Federated States of Micronesia, more than 75% of the matches with the WCPFC observers' data are under 2,000 days (less than 5 years) between the last record and the stranded position whereas matches with the PNA tracking data were less than 2 years. Similar patterns were found for Australia, New Caledonia, French Polynesia, and Wallis and Futuna (FIGURE 17).

Data from WCPFC and IATTC observer programs correspond to the last activity on the buoy that was recorded by observers, not the very last position recorded from a satellite buoy. Thus, it can overestimate the time difference between the actual last transmission and the stranding position, while fishers could potentially still have used the buoy and associated FAD, with this information not available to the observer. In addition, a floating object could have been stranded for a long period before being found by local communities. Consequently, the time difference can, again, be overestimated. Another point to note is that the PNA FAD tracking program started in 2016. No matching could therefore be found with buoys found stranded before 2016. Moreover, the PNA tracking database does not include the full trajectories of buoys, with some buoy trajectories having been "geo-fenced" (Escale et al., 2023) with the part of trajectories outside PNA country EEZs removed. Complete trajectories from both the WCPFC and the IATTC CAs would therefore be needed to identify more accurately the origin of buoys and the time buoys were drifting before reaching coastal areas. Note that high-resolution buoy data is not available for the IATTC until 2022 but is now mandatory for the whole fleet under Resolutions C-21-04 and C-24-01.



**FIGURE 17.** Boxplots of time difference between the date found stranded and the last known position in the fishery databases: the PNA FAD tracking data, the WCPFC observer data and the IATTC observer data (note that no confidential data was shared between organisations; e.g. vessel ID, flag ID, satellite buoy track); by PICT of stranding event. Grey dots indicate an individual stranding event. The lower and upper box boundaries indicate the 25<sup>th</sup> and 75<sup>th</sup> percentiles, respectively, the black line indicates the median, and the lower and upper error

lines indicate the 10<sup>th</sup> and 90<sup>th</sup> percentiles, respectively. AU = Australia; CK = Cook Islands; FM = Federated States of Micronesia; GU = Guam; HW = Hawai’i; NC = New Caledonia; PF = French Polynesia; PY = Palmyra; TO = Tonga; TV = Tuvalu; VU = Vanuatu; WF = Wallis and Futuna.

#### 4. Discussion and next steps

This paper presents the in-country data collection programmes related to dFADs found in coastal waters and on coastlines, as well as the development of a regional database. Data collection is now in place in 16 PICTs: American Samoa, Australia, Cook Islands, Federated States of Micronesia, French Polynesia, Guam, Hawai’i, Republic of the Marshall Islands, New Caledonia, Republic of Palau, Palmyra, Solomon Islands, Kingdom of Tonga, Tuvalu, Samoa, Wallis and Futuna. In the EPO, the voluntary collection of stranded FAD information has also been discussed, with some interested countries, and has started in Galapagos and potentially other EPO coastal states. In this paper, we identified that more than 3,500 stranding events have been identified, with data collected as far back as 2006. When compiling data from all the different programmes, considerable work in processing, and cleaning the data was necessary to standardise the data collected, while the database also includes links to supplied pictures that were sometimes used to complement the data entered. We also note that the data collection effort is spatially and temporally variable throughout the region, as most programs are based on voluntary reports from communities, while others used dedicated surveys (e.g., French Polynesia). In addition, in many places, the first step in the data collection effort was an inventory of all the buoys and FADs that had been stranded on the coastline. Future data collection will allow the identification of the rates of stranding events in a given time. It is important to highlight the growing involvement by more PICTs and the increased use of dedicated surveys, either with the use of drones, or directly in person, which improves temporal and/or spatial scope of stranding events in certain areas.

Data is currently collected at national level in each PICT on paper and later transferred to their own database through the use of Google Drive spreadsheets, and all of these individual databases feed a regional data collection effort held at SPC. With increasing reporting rates and number of PICTs

involved in the programme, the use of an app could be considered. In the future, dFAD stranding officers or coordinators may be needed to account for the large amount of data received.

As mentioned previously, some of the data collection programmes are independent initiatives in specific PICTs (

Appendix 2) another independent initiatives may also be occurring throughout the Pacific Ocean. In addition, in oceanic waters, some fishing companies likely sell or collect lost or abandoned dFADs from their own and other fleets and store them in port storage areas to be returned or traded back to company owners.

In a context where the data collection effort is variable throughout the Pacific region, discussions on the implementation of a voluntary-based data collection program on stranded FADs in the EPO have been initiated by the IATTC with their members with multiple CPCs showing interest to participate. IATTC and SPC staff have been collaborating in informal meetings to discuss elements needed to launch a voluntary regional data collection program in the EPO to harmonize and standardize data collection forms between the two organisations to the extent possible. The elements discussed included the need for developing regional awareness campaigns and communication materials, leveraging local initiatives, securing funding, and coordinating training sessions on data collection and data entry methods. Using a harmonized and dedicated form will facilitate data comparison and exchange between the two organisations, thereby strengthening collaboration and enhancing the regional data collection program to support Pacific-wide coverage, as recommended by the IATTC FAD-WG and the Commissions.

Relating to initiatives to reuse buoys found stranded throughout the Pacific and elsewhere, several buoy companies have launched their own repurposing program, such as Satlink “ReCon”, Marine Instruments “Blue Recovery” and Zunibal “Searchle” projects. SPC, on behalf and in partnership with some of its member countries and territories have joined these 3 initiatives. Satellite buoys used in the purse seine fishery have interesting functions such as GPS and echosounder which could equip artisanal anchored FADs and benefit local communities if they are found in good condition (e.g., Palau, New Caledonia, Cook Islands) or track marine debris (e.g. Australia, see

Appendix 2). While such devices remain expensive for many fisheries departments and organisations in PICTs, the option to give a second life to buoys found stranded present the double benefit of been beneficial for local communities and decrease the coastal pollution burden as there is often very limited options for local recycling. For buoys found broken, it is crucial to give recommendations on how to recycle and/or reuse different components and strengthen collaboration between buoy providers and fisheries departments and local associations working on this subject. As an example, Tangaroa Blue Foundation and Satlink are developing a recycling framework for the Satlink buoys, which involves designing a new tool to open up the buoy, as well as mapping recycling pathways for the components. Project ReCon partners will have access to these resources in the future.

In this paper, we presented an updated analysis on the data collected in the stranded FAD regional database (i.e., an update of FAD-07 INF-A). This highlighted the extent of FAD and buoy stranding events in the WCPO and some locations of the EPO. The type of stranding events, materials of the FADs found stranded, as well as the impacted habitats and the environmental damage detected were studied. Information collected through the data collection programmes and analysed here could also help prioritize and explore potential FAD retrieval programs in the future, as a measure to mitigate the impacts of lost FADs in the environment. Some PICTs such as French Polynesia, American Samoa or the Federated States of Micronesia manifested strong interests to launch a FAD watch/retrieval programme based on the FAD watch programme model implemented by TNC in Palmyra based on

collaboration with the fishing industry. A similar effort is being piloted in Galapagos by TUNACONS, and collected 48 FADs since 2022 in collaboration with local fishermen (FAD-09-RD-C).

A comparison with existing dFAD-related databases in the Pacific Ocean (e.g., WCPFC and IATTC observer data and PNA dFAD tracking data) was conducted and helped identify the origin (monitoring vessel, the flag and CA) and in many cases, some of the “life history” of the dFADs (area and date of last known position, drift patterns). Note that no confidential information was shared between organisations (e.g., vessel ID, flag ID, satellite buoy ID). Some buoys found stranded could not be matched with the fishery data investigated. This could be because; i) we did not have access to all the buoy trajectories in the Pacific Ocean (incomplete and modified trajectories from the PNA FAD tracking program in the WCPO and no confidential trajectory data for the IATTC CA at time of analyses); ii) observers cannot always record the buoy ID accurately from dFADs set on or visited by vessels; and iii) not all dFADs are set on or visited during their lifetime (see FAD-09-02 for a complete assessment of lifespan dynamics in the EPO). Under-estimation is likely with in-situ data as well, with not all the stranding events being reported or the identification number could be partially recorded. However, dissemination, communication and involvement of a large portion of the public, including fishermen and other stakeholders, but also the involvement of the fishing industry could help increase data collection and reporting levels and quality.

Additional countries and territories should consider implementing similar data collection programs and participating in this regional initiative. More reliable and relevant quantification of dFAD stranding or drifting nearshore, development of FAD recovery programs, as well as assessment of ecosystem impacts will be possible through long-term spatially-relevant data collection programmes, including countries and territories with low or no FAD fishing effort. Although the WCPFC and IATTC are currently moving forward in terms of fully non-entangling and biodegradable dFADs, such designs might still have an impact on the environment and sensitive habitats, making this programme relevant and timely. This could ultimately help inform dFAD management options in the Pacific Ocean.

### **Considerations and needs**

- Highlight the needs for and support of in-situ data collection and reporting programmes to better quantify dFAD stranding events and assess their impacts on marine and coastal ecosystems.
- Encourage additional PICTs, along with IATTC cooperating members and non-members (CPCs) to participate in the development, launch or enhancement of national data collection programs on stranded and lost dFADs, thereby strengthening regional collaboration, improving the database and contributing to Pacific-wide coverage;
- Acknowledge the need for FAD-buoy trajectory data, including historical data, from both the IATTC and WCPFC CAs, to better determine the origin of FADs and buoys, better determine the fate of FADs and to explore the development of spatial management strategies, recovery programs and incentive systems to reduce FAD loss, abandonment and stranding events;
- Highlight the need to develop initiative to reduce FAD loss and abandonment, including through potential offshore FAD retrieval programs;
- Encourage the development of other initiatives (e.g. FAD watch) and their participation by CPCs, to increase recovery of dFADs reaching coastal areas before they can cause environmental damages.
- Consider ways to develop solutions to process, re-use or recycle FAD materials and buoys in ports, and provide scientific-based advice to guide the management of dFADs in the Pacific Ocean.

## Acknowledgments

Funding for the awareness and data collection programs supported by SPC were provided by World Wildlife Fund (WWF), New Zealand Aid Program, Fond Pacifique, Trimarine and the World Bank. Mainui Tanetoa, Margot Boval, Jean-Claude Gaertner, Charles Daxboeck, Marie Soehnen, Anne-Marie Trinh are part of the data collection projects in French Polynesia, that are funded by the Direction des Ressources Marines, the University of French Polynesia/ Institut de Recherche pour le Développement (IRD) and The Nature Conservancy (TNC). Palmyra Atoll data collection is part of a larger dFAD Watch Program currently being developed and is funded by TNC and led by Kydd Pollock. Sarah-Jeanne Royer and Jennifer Lynch from the Center for Marine Debris Research at Hawai'i Pacific University are organising the data collection in Hawai'i. Tangaroa Blue Foundation would like to thank all of the Australian Marine Debris Initiative partners and volunteers involved in the collection and provision of data used in this report. For a list of AMDI partners who have contributed to the AMDI Database, please visit <https://www.tangaroablue.org>. The authors are grateful for the assistance provided by the fisheries department and officers in the field in each of the PICTs involved. The authors would also like to thank the local communities, fishermen, and general public involved in data collection in the PICTs. The authors are also grateful for any opportunistic report provided, including findings in Australia (Australian Fisheries Management Authorities); Cook Islands (Te Ipukarea Society); New Caledonia (Association Hô-üt); Pitcairn Islands (The Pew Charitable Trusts and The Plastic Odyssey); and Vanuatu. The authors would like to thank the members of the Parties to the Nauru Agreement, the Pacific Islands Regional Fisheries Observer Programme, and the Inter-American Tropical Tuna Commission for giving us access to their data for this analysis; as well as observers involved in the collection of observer data.

## References

- Balderson, S.D., Martin, L.E.C., 2015. Environmental impacts and causation of 'beached' Drifting Fish Aggregating Devices around Seychelles Islands: a preliminary report on data collected by Island Conservation Society. IOTC Technical Report IOTC-2015-WPEB11-39 15pp.
- Escalle, L., Hamer, P., PNA, 2023. Spatial and temporal description of drifting FAD use in the WCPO derived from analyses of the FAD tracking programmes and observer data | WCPFC Meetings. WCPFC-SC19-2023/EB-WP-05.
- Escalle, L., Mourot, J., Thellier, T., Lopez, J., Fuller, L.M., Royer, S.-J., Hood, L., Bigler, B., Jaugeon, B., Nicholas, T.-R.N., Pollock, K., Prioul, F., Marks, A., Jones, J., Lynch, J.M., Tait, H., Hamer, P., 2022. FAD-07-INF-A\_Analyses of the regional database of stranded (dFAD) in the EPO.
- Escalle, L., Scutt Phillips, J., Brownjohn, M., Brouwer, S., Sen Gupta, A., Van Sebille, E., Hampton, J., Pilling, G., 2019. Environmental versus operational drivers of drifting FAD beaching in the Western and Central Pacific Ocean. Scientific Reports 9, 14005. <https://doi.org/10.1038/s41598-019-50364-0>
- Zudaire, I., Santiago, J., Grande, M., Murua, H., Adam, P.-A., Nogués, P., Collier, T., Morgan, M., Khan, N., Baguette, F., Moron, J., Moniz, I., Herrera, M., 2018. FAD Watch: a collaborative initiative to minimize the impact of FADs in coastal ecosystems. IOTC Technical Report IOTC-2018-WPEB14-12 21pp.

## Appendices

- Appendix 1. Poster depicting the data collection program for the Cook Islands in English
- 
- Appendix 2. Description of stranded FAD data collection in independent **programs**
- Appendix 3. Data collection form for fisheries officer

- Appendix 4. Description of data collection for FADs found stranded or at sea
- Appendix 5. Supplementary figures

APPENDIX 1. Poster depicting the data collection program for the Cook Islands in English

# FISH AGGREGATING DEVICE (FAD)

## DRIFTING FAD FOUND BEACHED OR AT SEA ?

Where is the buoy ID number ?



ISL+123456   DSL+123456   MSI+123456   T7+123456789   P1234NF   123456  
or Ze0123456789



**WHAT IS A DRIFTING FAD?**

It is a raft, generally made of bamboo, with a tail of net, cords and/or canvas and a satellite buoy, deployed by tuna purse seiners to aggregate and catch tuna.

**WHY ARE WE COLLECTING THIS DATA?**

To quantify the number of beached or lost FADs, and to note their impact on coastal areas, which will help improve the management of FAD fishing.

**WHAT TO DO WITH THE FAD?**

If possible, tow the drifting FAD back to shore then contact the MMR office.

**RECORD ANY OF THESE DETAILS:**

- What did you find ?
  - a FAD by itself
  - a FAD with a buoy
  - a buoy by itself
- Buoy ID number and any mark painted on the buoy
- Date found
- Location (Lat/Lon or name of beach, village, island...)

**IF POSSIBLE, NOTE:**

- Environment: at-sea, coral reef, beach, lagoon
- Materials: bamboo, net, cord, floats
- Tail length (if possible)
- What did you do with the FAD/buoy? (e.g. removed from water or land, left drifting, sunk, fished)
- Any additional comments? (e.g. environmental damage, entangled animals or aggregated tuna or other animals)

**TAKE PICTURES:**

- General picture of what you found
- A close-up of the buoy with the ID number visible

**SEND AN EMAIL TO:** [rar@mmr.gov.ck](mailto:rar@mmr.gov.ck)

**OR CALL**                    **28721**



Pacific Community  
Communauté du Pacifique



Ministry of Marine Resources  
Ministère des Ressources Maritimes



WWF

© IGC 309 – Illustration: Ben Cole

## **APPENDIX 2. Description of stranded FAD data collection in independent programs**

At Palmyra Atoll, TNC and the U.S. Fish and Wildlife Service (USFWS) have collected data on dFAD strandings since 2009. Visual surveys across shallow reefs, lagoon flats, and beaches have been opportunistically linked with other projects; however, now that consistent stranding areas have been established, specific surveys are being scheduled across all 12 months of the year. Designs of the stranded dFADs, the materials used, and the environmental impacts are described. When a satellite buoy is present, and the identification number is visible, it is also recorded. A dFAD Watch-type program (Zudaire et al., 2018) has also been in place at Palmyra Atoll since June 2021. In this program, fishing companies alert local partners if a dFAD comes close to Palmyra Atoll's shores, so it can be removed before causing any environmental damage.

In Hawai'i, the stranded dFAD collection was first initiated by Sarah-Jeanne Royer as a member of Nikolai Maximenko's group at the University of Hawai'i at the International Pacific Research Center. The program is now being monitored by the Center for Marine Debris Research (CMDR) at Hawai'i Pacific University (HPU). The data collection started in 2014 and has expanded to include several collaborators that report the findings to the research group. When the geographical location of the dFADs is known, some buoys are re-directed to the island of Oah'u and stored in a warehouse to potentially repurpose the buoys to tag and track marine debris like fishing nets. A recent project provides a financial incentive to commercial fishermen that retrieve derelict dFADs at sea, which has resulted in the Hawaiian longline vessels removing several complete dFADs from fishing grounds far offshore of Hawai'i.

French Polynesia has also started a large project to quantify the number of dFADs drifting within its EEZ, including the number of stranded dFADs, and their ecosystem impacts. The program involves several components: i) data reported by local communities through a form that can be directly downloaded or filled in on the marine resources authority's website (<http://www.ressources-marines.gov.pf/dcpech>); ii) dedicated surveys in 9 islands of the Tuamotu (Hao, Amanu, Raroia, Rangiroa, Reao, Tikehau, Tureia, Raraka, Fakarava), with visits to local communities, shoreline surveys using a drone, shore cleanings, and FAD recycling operations.

In Australia, Tangaroa Blue Foundation (TBF) coordinates the Australian Marine Debris Initiative® (AMDI), an on-ground network of volunteers, communities, organisations and agencies around the country removing, documenting, and preventing marine debris and plastic pollution. The AMDI Database is the largest marine debris database in the southern hemisphere, with more than 23 million litter items recorded at more than 4,300 clean-up sites since 2004. Marine debris data are collected via community clean-ups or as part of regular site-specific monitoring programs. In particular, data on dFAD strandings have been recorded in the AMDI Database since 2004 across Australia's coasts, with the majority found along the coast of Queensland. Satellite buoys were recorded frequently prompting Satlink, one of the buoy providers, to partner with TBF to develop Project ReCon: a recover, repair, reuse and recycle program of satellite buoys. TBF and their partners in the AMDI collect satellite buoys found during clean-up events and Satlink then liaises with the industry to facilitate reassigning ownership of these buoys from the commercial fishing fleet to TBF. Once a buoy is part of the project ReCon, Satlink and TBF check the buoy's condition and find a suitable re-use project, such as scientific research, tagging and recovering ghost nets, etc. Buoys are then stored by community partners from a variety of sectors (i.e., tourism, charter operators, and Aboriginal and Torres Strait Island Rangers), so that they can be deployed on ghost nets that cannot be immediately recovered when located due to their size or the capacity of the vessel that found them. Previously, recordings of dFADs and buoys were limited to stranding events; however, Project ReCon helps improve understanding of dFADs found in the coastal waters of Australia. Historical data that were transferred to the regional database at SPC focused mainly on satellite buoys, however, historical information related to dFADs exist and will be transferred to the SPC database in the future.

In Galapagos, a program led by Galapagos Conservation Trust and the Galapagos National Park has been in development since 2024, with data that has started to be collected, but not share with SPC or IATTC yet. The programme will aim to evaluate both opportunistic sightings of dFADs and pilot data where dFADs have been quantified along remote coastlines using drone surveys and during coastal clean-up activities. In addition to these efforts, private initiatives such as a voluntary dFAD recovery program started in 2022 by the Tuna Conservation Group (TUNACONS) – a consortium of Ecuadorian tuna fishing companies, has recovered dFADs thorough coordination between the TUNACONS – adhered fleet, local coordinators and local fishers (see FAD-09-RD-C).



# FAD Sighting form v3

**Form details** Date: \_\_\_\_\_ Form nb: \_\_\_\_\_  
 Completed by: \_\_\_\_\_

**2/2**

**Raft materials** (Tick one or several)

**Raft materials structure and flotation:**  Bamboo  Log  PVC  Floats  Plastic drum  Fiberglass drum  Metal drum  
 Steel  Polystyrene  Unknown  Other: \_\_\_\_\_

**Raft materials covering:**  None  Ropes  Nets  Plastic sheeting  Canvas  Unknown  Other: \_\_\_\_\_

**If net present in the raft, mesh size :**  Small (<7cm)  Large(>7cm)  Small & Large  Unknown

**Estimated size of the raft (m) (Length x Width):** \_\_\_\_\_ x \_\_\_\_\_ or  Unknown

**Underwater component/tail** (Tick one or several)

**Submerged tail presence (i.e., part of the FAD normally under water):**  Yes  No  Unknown

**Submerged tail materials:**  Unknown  Net  Rope  Canvas  Plastic sheeting  Bamboo  Fishing lines  
 Other: \_\_\_\_\_

**Design of the tail:**  Open net, mesh size:  Small (<7cm)  Large(>7cm)  Other: \_\_\_\_\_  
 Net rolled up in bundle, mesh size:  Small (<7cm)  Large(>7cm)  Other: \_\_\_\_\_  
 Other: \_\_\_\_\_  
 Unknown

**Estimated depth of submerged tail (m):** \_\_\_\_\_ or  Unknown

**Fate of the FAD**

**Fate of the FAD?**  Left in the environment  Sunk  Raft removed, tail section left  Unknown  Removed from the environment (tick if "found in a private property")  Other: \_\_\_\_\_

**Only if removed from environment, purpose:**  Burned  Left with the finder  Landfill  Research  Recycled  Re-used (specify): \_\_\_\_\_  Other: \_\_\_\_\_  Unknown

**Impact on / interaction with marine life** (Tick one or several)

**Environmental damages caused by the FAD :**

Entangled animals:  Yes  No  Unknown

Entangled on corals:  Yes  No  Unknown

**Entangled animals?**  Turtle  Shark  Fish  
 Marine mammal  Unknown  Other: \_\_\_\_\_  
**Status:**  Dead  Alive  Unknown  
**Species (if known):** \_\_\_\_\_  
**Number of individuals:** \_\_\_\_\_

**If FAD is entangled on coral reef, please state the approximate size of the area impacted (m<sup>2</sup>):** \_\_\_\_\_

**Fish caught during a set on the FAD:**  No  Yes  Unknown

If yes, **Species (if known):** \_\_\_\_\_

If yes, **Weight of the catch (in kg) (if known):** \_\_\_\_\_

**Number of individuals:** \_\_\_\_\_

**Fish or other animals aggregated around the FAD :**

No  Yes  Unknown

If yes, **Species (if known):** \_\_\_\_\_

**Number of individuals:** \_\_\_\_\_

**Comments:** \_\_\_\_\_  
 \_\_\_\_\_

**Number of pictures:** \_\_\_\_\_

## APPENDIX 4. Description of data collection for FADs found stranded or at sea

### Description of data collection for FADs found stranded or at sea

For further informations, contact Jennifer Mourot [jenniferm@spc.int](mailto:jenniferm@spc.int)  
or Lauriane Escalle [laurianee@spc.int](mailto:laurianee@spc.int)

#### **Why are we collecting these data?**

We are collecting these data in order to quantify the number of lost and stranded FADs, and to note their impact on coastal areas, which will help improve the management of FAD fishing. Drifting FADs are always deployed with a satellite buoy, so that fishers know the position of their FADs. This buoy is usually also equipped with an echosounder to estimate the amount of tuna aggregated underneath. Fishing companies have started sharing data both of the FAD's position, as well as the echosounder data from the satellite buoys deployed on FADs. These data are used in scientific studies that guide management of FAD fishing. When FADs are found at sea or stranded, it is therefore very important to record the unique buoy ID number, to potentially match found FADs with these existing datasets.

However, fishers commonly remotely deactivate satellite buoys when FADs drift outside fishing areas. The dataset transmitted by fishing companies hence only gives a partial image of the FAD trajectories, and the number of stranding events is underestimated in this dataset. Therefore, having access to additional information on stranding events, but also on FADs drifting in coastal areas (with the buoy ID number, if still attached to the FAD), will help complement the existing dataset and better estimate the impact that FADs may have on coastal areas.

#### **Description of the fields in the spreadsheet**

- **Entry number**  
(Internal use only. Number of the object found (1 to n). Used to rename the pictures.)
- **Entry number from independent program**  
For independent programmes that already have their own numbering.
- **Entered by**  
Name of the person entering the data.
- **Date entered**  
Date of data entry. Use the yyyy/mm/dd date format.
- **Type of data**  
Specify how the data was collected, was it a dedicated survey, using a drone or by people (e.g. walking along a coastline), opportunistic reporting (not part of an existing programme) or by local communities (part of a data collection programme).
- **Found by**  
Name of the person who found the FAD and/or the satellite buoy.
- **Contact**  
Enter contact detail (email address, phone number) of the person who found the FAD and/or the satellite buoy.
- **Date found (yyyy/mm/dd)**  
Date when the object has been found. Use the yyyy/mm/dd date format.

However, if the object was previously collected by local communities some time ago (for example, stored in a garden or at the port), record here the date when the officer in charge of the data collect, found or saw this object; then enter the date when the object was actually found in the "Initial date found by communities" field (see below).

- **Island**

Note only the name of the island where the object has been found.

- **Location**

In particular if the lat/lon were not recorded, record where the object was found, e.g. name of beach, town, island, etc.

- **Accurate GPS coordinates**

Specify if the exact GPS coordinates where the floating object was found are available.

- **Latitude and longitude (If provided)**

Record latitude and longitude in decimal format.

- **Environment (if provided or visible on the pictures)**

Where the FAD and/or buoy has been found: drifting at-sea in the lagoon or the ocean, on a beach, a coral reef, a rocky shore, a mangrove; or previously found and reported from a garden, a wharf (if the object has been found some time ago), etc.

- **Initial date found by communities (optional)**

When reports relates to floating objects already collected previously by communities, record the date it was originally found if it is known. If the precise date is unknown, it could be approximated. Use the yyyy/mm/dd , yyyy/mm or yyyy date format.

- **Initial location or coordinates (optional)**

When reports relates to floating objects already collected previously by communities, record the location/coordinates where it was originally found, if they are known.

- **Initial environment when found by communities (optional)**

When reports relates to floating objects already collected previously by communities, record the environment it was originally found, if it is known.

- **Buoy present**

Was a satellite buoy present (i.e. buoy attached to a FAD or buoy by itself)? Yes/No.

- **Buoy type**

What kind of buoy was found? Most of the buoys found with FADs are satellite buoys (whole or part of it), but it could also be oceanographic buoy or radio buoy (used by longline vessels for instance). If you are not sure, take photos or look at the photos below to orientate the choice.

Satellite buoy	Radio buoy	Oceanographic buoy
		 

- **Buoy ID number (very important if a buoy is present)**

Enter the satellite buoy ID number (see the end of this document how to find the Buoy ID number depending on the buoy brand).

Examples of satellite buoy ID number:

**DL**+123456 ; **ISL**+123456 ; **DSL**+123456 ; **SLX**+123456

**M3I**123456; **M3**+123456; **M4**+123456

**T07**123456789; **Te7**123456789; **T7**+123456789; **T8X**123456; **F8E**123456789 ; **Z07**123456789

**P1234NF**; **P1234N**; **WF1234N**; **CN123N**

123456

- **Buoy brand**

Record the brand of the buoy, usually written on the top or the side of the buoy. Most brand used are Satlink, Marine Instruments, Zunibal, Kato, and Ryokusei.

- **Buoy model**

After identifying the brand of the buoy, record the model. Either another name is written inside or on the side of the buoy, or record the letters at the beginning of the buoy ID number. Examples of buoy ID number, with the buoy model in bold:

**ISL**+123456 ; **DSL**+123456 ; **SLX**+123456 : These types of letters correspond to a Satlink buoy (non-exhaustive list).

**M3I**123456; **M3**+123456; **M4**+123456 : These types of letters correspond to a Marine Instruments buoy (non-exhaustive list).

**T07**123456789 ; **Te7**123456789 ; **T7+**123456789 ; **T8X**123456 ; **F8E**123456789 : These types of letters correspond to a Zunibal buoy (non-exhaustive list).

- **Buoy condition**

Note the condition of the buoy. Determines if the buoy is whole or on parts. If it is whole, note any damage such as cracks on the top and/or bottom plastic case, if there is water inside, or if the echosounder is cracked. If the buoy found is not complete, what part did you find? It could be the plastic case (top and/or bottom), the electronics or both.

- **Marks on the buoy (if provided or visible on the pictures)**

Record any marks painted on the satellite buoy. Could be a vessel name, or the abbreviation of a vessel names, just a letter, a number, a number and a letter, and sometimes the buoy ID number.

- **Fate of the buoy**

What has been done with the satellite buoy: removed from the environment, left drifting, left on shore, sunk, etc.

- **Purpose if buoy removed**

If the buoy has been removed from the location it was found, mention why it has been removed: for research, avoid pollution, recycling (use battery, solar panels...), etc.

- **Buoy storage location**

If the buoy has been removed from the location it was found, record where the buoy is stored (if known). Particularly important information in the context of the project of reuse of old satellite buoys.

- **FAD present**

Was a FAD present (i.e. FAD by itself or FAD with a buoy)? Yes/No.

- **FAD Type**

Note if the object found was a drifting FAD (dFAD), an anchored FAD (aFAD; industrial aFADs commonly look like a giant drum) or just a part of a drifting FAD.

Examples of a drifting FAD, drifting FAD parts, and an anchored FAD.

Drifting FAD	Drifting FAD parts	Anchored FAD
		

- **FAD condition (if provided or visible on the pictures)**

What is the condition of the FAD when found? Intact, beginning to break, mostly fallen apart.

- **Marks on the FAD (if provided or visible on the pictures)**  
Record any mark attached to the FAD or painted on it.
- **Raft shape**  
Define the shape of the FAD. Generally, FADs have a rectangular or a square shape, eventually hexagonal, but sometimes they look like a sausage of buoys wrapped in netting.
- **Raft materials – Structure and flotation (if provided or visible on the pictures, can be multiple entries)**  
List all the materials making the raft of the FAD, especially materials for the structure and the flotation: bamboo, log, floats, plastic or metal drum, polystyrene, etc.
- **Raft materials – Covering and other (if provided or visible on the pictures, can be multiple entries)**  
List all the materials making the raft of the FAD, especially covering materials or other types of materials: net, rope, canvas, plastic sheetings, etc.
- **If net is present (raft), mesh size**  
Determine the mesh size of the net used for the raft, if it is small (under 7 cm) large (more than 7 cm), or both types of netting are present.
- **Length and Width of the raft (if estimated)**  
Estimate the size of the FAD raft, Length (m) and Width (m).
- **Tail presence**  
Is there any submerged appendage under the raft structure? Yes/No.
- **Tails materials (if provided or visible on the pictures, can be multiple entries)**  
List all the materials making the tail of the FAD (underwater appendages): bamboo, log, net, cord, canvas, etc
- **If net is present (tail), mesh size and design**  
Determine the mesh size of the net used for the tail, if it is small (under 7 cm), large (more than 7 cm) or both types of nets are present. Determine as well, the design of the tail. It could be a open netting panel or netting could be rolled up into a bundle.
- **Tail length (if estimated)**  
Estimate the length of the FAD tail, i.e. the materials (rope, net, etc.) hanging under the FAD raft, in meters.
- **Fate of the FAD (if provided)**  
What has been done with the FAD: removed from the environment, left drifting, left on shore, sunk, fished, etc.
- **Purpose if FAD removed**  
If the FAD has been removed from the environment it was found, mention why it has been removed: for research, landfill, burned, stored (to do what?), recycled (to do what?), etc.
- **Environmental damage (if provided or visible on the pictures)**  
Any environmental damage recorded: e.g. corals and/or animals are entangled in the tail of the FAD.

- **If FAD is entangled on coral reef, please state the approximate size of the area impacted (m2)**

- **Entangled animals in the FAD ? (if provided or visible on the pictures)**

**(Yes/No Animals Species Dead/Alive Number of individual)**

Record if any animals were found entangled on the net beneath the FAD and/or the net used to cover the raft. If possible, record the type of animal, the species (if known), if the animal is alive or dead, and the number of individuals.

- **Aggregated fish or animals ? (if provided)**

**(Yes/No Animals Species Number of individual)**

Record if any fish or other animals were seen aggregated under the FAD. If possible, record the type of animal, the species (if known) and the number of individuals.

- **Fish caught ?**

**(Yes/No Animals Species Weight Number of individual)**

Record if any fishing was performed around the FAD. If it was the case, mention the species (if known), the number and/or the catch in kg.

- **Other comments**

Any other comments: e.g. some tunas were aggregated under the FAD, the FAD could not be removed because too heavy, materials reused as fishing gear, etc.

- **Number of pictures received**

Record how many pictures have been received/taken. If no pictures exist, put 0. If some pictures exist but you are waiting for their transfer, put "waiting for photos".

- **Pictures name**

Rename the pictures using a unique identifier containing, country, entry number (first field) and date found (if date found is unknown use the date entered).

<CountryCode>\_<Seq. No.>\_<Date:YYYYMMDD> Ex: FM\_1\_20190923

Add another number if more than one picture: e.g. FM\_1\_20190923\_P1; FM\_1\_20190923\_P2; FM\_1\_20190923\_P3. Then copy the pictures in the folder in google drive.

- **Buoy ID number verified**

Has the satellite buoy ID number been verified by the fishery officer on a picture or directly: Yes/No.

- **SPC Check**

For SPC staff only, check information related to the entry and compare with photos (if available).

- If photos are available and there are no questions, the entry's check will be marked as "Yes".

- If there are questions regarding entry by SPC staff, it will be marked as "Yes, but waiting for further information".

- If there are no photos although it is recorded that they exist, and the line has been checked, it will be marked as "Yes, but waiting for photos".

- If the line has been checked and there are no existing photos, it will be marked as “Yes, but nothing to check”.

- If the line has not been checked, it will be marked as “No”.

- **If dedicated survey, survey name**

If it was during a survey, note the name as the following format:

<country acronym>\_CS\_<date of the survey in an YYYY\_MM\_DD format>

Example: a survey in Tonga the 15<sup>th</sup> of february 2023 : TO\_CS\_2023\_02\_15

If there are several surveys the same day, please add a “P” (for “part”) and a number at the end:

Example: TO\_CS\_2023\_02\_15\_P1 ; TO\_CS\_2023\_02\_15\_P2 ; TO\_CS\_2023\_02\_15\_P3 ...

- **Length of the area surveyed (m)**

If it was during a survey note the total length surveyed in meters.

- **If dedicated survey, total area (m2)**

If it was during a survey note the total area surveyed in squared meters.

- **If dedicated survey, reef area (m2)**

If it was during a survey note the total reef area surveyed in squared meters.


- **If dedicated survey, lagoon area (m2)**

If it was during a survey note the total lagoon area surveyed in squared meters.


**Information on satellite buoys**

Buoy brand	Special features	ID number
<p><b>Marine Instruments</b></p> 	<ul style="list-style-type: none"> <li>• Present big black handles on sides</li> <li>• Often have a green bottom plastic case.</li> <li>• Top plastic case is higher than other buoys brand.</li> </ul>	<p>The ID Number is present on the side of the buoy and the top plastic case at the corners, or on a small metal plate inside between the solar panels.</p> 


<p><b>Satlink</b></p> 	<p>Often has 6 solar panels radiating from the center (looks like an asterisk/star or flower)</p>	<p>In the center of the buoy inside a (black) circle.</p> 
<p><b>Zunibal</b></p> 	<ul style="list-style-type: none"> <li>• Often 4 cross-shaped solar panels with a blue background.</li> <li>• The top plastic case is quite flat.</li> </ul>	<p>The model of the buoy (here at the top) and the number (here at the bottom) is in the center of the buoy.</p> 
<p><b>Kato</b></p> 	<ul style="list-style-type: none"> <li>• Often 4 solar panels arranged in a square</li> <li>• Dark green background</li> <li>• Fairly flat top plastic case</li> </ul>	<p>It is located on a metal plate at the ends of the solar panels. The code on the first line is the model of the buoy, the code below is the ID number.</p>

		
<p><b>Ryokusei</b></p> 	<ul style="list-style-type: none"> <li>• Few information available.</li> <li>• Often 2 solar panels side by side</li> <li>• White background.</li> </ul>	<p>The ID number is on a small white box inside.</p> 


APPENDIX 5. Supplementary figures



**FIGURE S5.1.** Map of stranding events by type of object in some islands of the Tuamotu Archipelago (top) and Marquesas Islands (bottom) (French Polynesia) between 2019–2025. Islands inside orange squares are islands where dedicated surveys using drones occurred.



**FIGURE S5.2.** Map of stranding events in Rarotonga (left) and in Aitutaki (right) (Cook Islands) by type of object between 2018–2025.



**FIGURE S5.3.** Map of stranding events in Australia by type of object between 2018–2025.



**FIGURE S5.4.** Map of stranding events in Wallis (left) and Futuna (right) by type of object between 2018–2025.